

Health & Hygiene

6th Term I Unit 6 – Health & Hygiene

- As defined by World Health Organization (WHO), it is a "State of complete physical, mental, and social well being, and not merely the absence of disease or infirmity." Health is a dynamic condition resulting from a body's constant adjustment and adaptation in response to stresses and changes in the environment for maintaining an inner equilibrium called homeostasis.
- Hygiene is a science of the establishment and maintenance of health conditions or practices (as of cleanliness) conducive to health has poor personal hygiene. Brushing your teeth regularly is an important part of good oral hygiene. Hygiene is the practice of keeping yourself and your surroundings clean, especially in order to prevent illness or the spread of diseases.

Components of Food

- The Chemical constituents of food which give us energy, help to build our body and protect us from diseases are called Nutrients.
- Carbohydrate
- > Proteins
- ➤ Fats
- Vitamins
- Minerals



➤ Water

Carbohydrates

Carbohydrates are energy giving component of the food.

S.No	Form of Carbohydrates	Sources
1	Sugar	Fruits, Honey, Cane Sugar, Sugar Beet
2	Starch	Rice, Wheat, Maize, Potato, etc.
3	Dietary fibre	Whole grain, nuts, etc.

Fats

• Fat is also an energy-giving food and provides more energy than Carbohydrates. Some important sources of fats are butter, ghee, milk, cheese, paneer, nuts, meat, fish, egg yolk etc. Apart from giving energy, they insulate our body and protect the cells

PROTEIN

Body Building Foods

• Proteins are necessary for our growth and repair, as well as for regulating various body functions such as digestion. The sources of proteins are pulses, eggs, fish, milk, chicken, soya bean, nut, grams etc, Proteins are body building foods.

ENTRE

Soyabean is the highly rich source of protein

Vitamins

• Vitamins are required for carrying out various biochemical reactions in our body. Fruits, vegetables, grains, meat products are good sources of vitamins. Vitamins are called as protective food. There are six major vitamins A, B, C, D, E and K. Vitamins B and Vitamins C are water soluble, Vitamins A, D, E and K are fat soluble.



Vitamin	Found	Disease we get	Symptoms
	abundantly in	if deficient in	
		this	
Vitamin A	Fish oil, egg,	Night blindness	Poor vision,
	milk, ghee,		difficulty in
	carrot, corn,		seeing in dim
	Yellow fruits,		light
	greens	D 11 1	
Vitamin B	Whole grain,	Beri beri	Nervous
	unpolished rice,		weakness,
	milk, fish, meat,		fatigue.
	peans, lentils Green		
	Vegetables		
Vitamin C	Oranges,	Scurvy	Bleeding gums
v italiili C	Gooseberry,	Scurvy	biccuing guins
	Greenchilly,		
	Tomato		E
Vitamin D	Fish oil, milk	Rickets	Weak, flexible
	and eggs. It is	TEN	bones.
	also made in our	LL	
	skin using		
-	sunlight		
Vitamin E 💦 🔊	Vegetable oils,	Nervous	Childlessness,
	Green	weakness,	lack of resistance
	Vegetables,	dimming of	1
	whole wheat,	eyesight	illnesses
	Mango, apple,		
X7: . X	greens	TAT 1 C (1	T 11 (
Vitamin K	Green	Weakness of the	Even a small cut
	Vegetables,	bones, teeth etc.	bleeds profusely.
	Tomato,		
	cabbage, eggs,		
	milk products.		

Gooseberries contains nearly 20 times the vitamin C than Orange



Minerals

• Minerals are required for growth as well as for the regulation of normal body function.Green leafy vegetables like spinach, pulses, eggs, milk, fish and fruits are important sources of minerals in our diet. Minerals are also a protective foods.

Minerals	Functions
Calcium	Strong bones and teeth, clotting of blood
Phosphorus	Strong bones and teeth
Iodine	Synthesis of thyroid hor- mone
Iron	Formation of haemoglobin and brain development

80% of the world production of Moringa Leaves is in India. The Major countries which import Moringa Leaves are China, US, Germany, Canada, South Korea and European countries

Water

• Our body needs an adequate supply of water is order to maintain good health. Any human being should take minimum eight tumblers (2 Litres) of water every day.

Health

- Health is a state of complete physical, mental and social well-being and not merely absence of diseases. Eating a healthy diet keeps you physically and mentally fit. When you are physically healthy, you feel confident you are more outgoing and have a greater capacity for enjoying life.
- Unhealthy food choices lead to obesity and illness, preventing you from socializing with friends and family. So choose your diet carefully.



Balanced Diet

- A diet should contain adequate amount of all the necessary nutrients required for healthy growth and activity.
- An increased capacity to work
- Good physical and mental health
- Increased capacity to resist diseases.
- Help in proper growth of the body.
- A balanced diet contains sufficient amount of various nutrients to ensure good health. Food should also provide the appropriate amount of energy and adequate amount of water.

Malnutrition:

• Malnutrition occurs when all the nutrients that the body needs are not obtained in the proper proportions from the diet. The word malnutrition refers to the condition that results when a person does not take a balanced diet. Malnutrition leads to deficiency disease. The diseases that are caused due to lack of Nutrients in the diet are called Deficiency Diseases.

India has the second highest number of obese children in the world after China, according to a study that has found that 14.4 million children in the country have excess weight.

Protein and Mineral Deficiency Diseases

Protein Diseases	Symptoms		
Kwashiorkar	Stunted growth, swelling of face		
	and limbs, Diarrhoea.		
Marasmus	Skinny appearance, slow body		
	growth.		

Mineral	Deficiency Disease		
Calicum	Rickets		
Phosphorus	Osteomalatia		
Iodine	Cretinism (in child) Goitre (in		

		CHENNAL
	adult)	
Iron	Anaemia	

Physical Exercise and Rest

- Physical exercise is any bodily activity that enhances or maintains physical fitness and overall health and wellness. It is performed for various reasons, including
- increase in growth and development,
- strengthening muscles and the cardiovascular system,
- developing athletic skills, weight loss or maintenance, and enjoyment.
- Physical exercise may help to decrease some of the effects of childhood and adult obesity.

Deep sleep seems to be one of the most critical time for body repair

REST

• Proper amount of rest is essential for physical and mental health. Rest is as important as nutrition and physical activity for growth and development and good health.

Personal Cleanliness

- Hygiene is a set of practices performed to preserve health. According to the World Health Organization (WHO), "Hygiene refers to conditions and practices that help to maintain health and prevent the spread of diseases."
- **Personal hygiene** involves those practices performed by an individual to care for one's bodily health and well being, through cleanliness. It includes such personal habit choices as how frequently to bathe, wash hands, trim fingernails, and change clothing. It also includes attention to keep surfaces in the home and workplace, including bathroom facilities, clean and pathogen-free.



Introduction of Microbes

- The diseases or conditions caused by microorganism due to the negligence of personal hygiene.
- ➢ Diarrhoea
- ➤ Tooth decay
- Athlete's foot(Madurai's foot)
- ➤ Dandruff.

Most of the microbes belong to four major groups:

- ➢ Bacteria
- ➢ Virus
- Protozoa
- ➤ Fungi

Bacteria

- Bacteria are very small prokaryotic microorganisms.
- Bacterial cells do not have nucleus and do not usually have membrane bound organelles.
- > Bacteria can exist either as independent organisms or as parasites
- They invade tissues
- They produce pus or harmful wastes

Bacterial Diseases

S.No	Bacterial diseases	Mode of transmission
1	Cholera	Contaminated water
2	Pneumonia	Inhalation of airborne droplets
		from a sneeze or cough
3	Tetanus	Contamination of wounds with
		the bacteria.
4	Tuberculosis	Inhalation of airborne droplets
		from sneeze or cough
5	Typhoid	Contaminated food or water



Disease

Disease is a definite pathological process having a characteristic set of signs and symptoms.

Disorder

Disorder is a derangement or abnormality of function.

Virus

• Virus is an acellular agent that replicates only inside the cells of other living organisms. Virus can infect all types of life forms plant, animals and microorganisms. They invade living normal cells and use their cell machinery to multiply. They can kill damage or change the cells and make you sick.

GENTR

Diseases Caused By Virus

- Common cold
- ≻ Influenza
- > Hepatitis
- > Polio
- ➤ Smallpox
- Chicken pox
- ➤ Measles

Microscopes help to study the structure of the microorganisms



7th Term I

Unit 6 – Health & Hygiene

- Health is the best wealth. If you have good health, you will have a sound mind and you will gain good knowledge and wealth also. To maintain good health, you should follow good hygiene, eat nutritious food, do exercise, take rest and have a sound sleep.
- It is also refers to a state of a sound mind and body free from any sickness or ailment, stress and problems. In simple words, health refers to the physical, emotional and psychological well-being of a person.
- Hygiene refers to the good habits and their practices which is followed to prevent diseases, maintain good health, especially through cleanliness, consumption of safe drinking water and proper disposal of sewage. It refers to all those activities that are done for improving and maintaining good health and sound mind.

Cleanliness

- Cleanliness refers to the maintenance of personal and environmental hygiene. In simple words, It refers to the state of being clean which is essential for good health. To protect us from diseases it is essential to maintain good health by taking regular bath, cleaning the clothes and surroundings and also avoiding unhygienic food consumption.
- Personal hygiene is defined as "the branch of health which is concerned with the individual's adjustment to the physiological needs of the body and mind for the attainment of the maximum level of health, it also refers to the cleaning and grooming of the body.
- Colds and the flu are common communicable diseases. It is caused not only by bacteria but also by virus. When you have cold and flu, you may also have running nose, cough, sore throat, and sometimes



fever or pain in the joints. For some, this condition may also lead to mild diarrhoea.

Secretions oozing out from the nose may contains the bacteria or virus. When the patient touch the nose and some other object or someone else the virus is transferred. When the patient sneezes or coughs the virus comes out with the droplets and become airborne. Hence it is a good practice for the patient with cold and flu to use a handkerchief to blow the noses and also wash the hands often to ensure that they do not accidentally spread the virus to others.

Community Hygiene

- A community is formed by a group of people living together in a particular area. If the people in a community wish to lead a healthy life, they should maintain basic community hygiene. It can be done by adopting the following measures.

- Drains should be covered properly.
 The domastic The domestic wastes should be segregated and properly disposed off safely in separate dust bins provided by the Government (Green and Blue).
- Used water from houses should not be let out into open drains or open areas

Dengue is spread by mosquitoes of Aedes aegypti caused by DEN-1, 2 virus belonging to the type- flavivirus. It decrease counting of the blood platelets of human blood and it has a maximum flight range of 50-100 meters in and around the places.

Care of the body

• A human body is a massive miracle. It consists of organs and systems, which functions day in and out. Our body in compared to a machine. The human body systems work well with proper maintenance and guidance. For smooth functioning, all the parts of



the body should work in unison. The digestive system, circulatory and muscular system is the core systems that should be in synchronization and functioning well. So keep them well by proper care.

Dental Care

- Dental care or broadly speaking oral hygiene is an important aspect of the personal health of an individual. Good oral hygiene implies sound teeth and healthy gums with healthy surrounding tissues. The physical act of chewing food promotes saliva and gastric secretions which helps digestion. The act of chewing and tasting is called 'mastication'. It gives pleasure and emotional satisfaction of eating food. Teeth is essential for good appearance and clear speech also.
- When you brush two times a day, it will prevent the formation of tartar and plaque on your teeth and gums.
- When you Floss, it will remove food particles, plaque, and bacteria which build up between your teeth. When you start flossing, your gums may bleed a little bit, but after few days that will be stopped. It should be started only with proper medical guidance.

Diseases affecting the teeth

• Diseases affecting the teeth and gums, their causative agents and remedial measure are given below:

S.No	Name of the	Causative	Impacts/	Remedial
	Diseases	Agents	Consequences	measures
1	Bleeding	Vitamin C	Bleeding of	Eating citrus
	Gums	deficiency	the gums	fruits.
2	Tooth decay	Bacteria in	Bacteria	Brushing
		plaque	produce acids	and flossing
				the teeth can
				prevent
				decay.
3	Periodontitis	Tobacco	Severe form of	Chewing
		chewing	gum disease	type of

		(CHENNAI
	ruin	the	tobacco
	bones,	gums,	should be avoided.
	and	other	avoided.
	tissues.		Eat a well -
			balanced
			diet.

Eye Care

• Eyes are an important organ of our body. They are considered as "windows to the world". Eyesight is the most important sense. 80% of what we perceive comes through the sense of sight. The protection of the eyes, can reduce the odds of blindness and vision loss. we protection of our eye from the diseases, surroundings, climate condition.

Diseases affecting Eye

Disease affecting the eye and their remedial measure are given below:

Hair Care

• The condition of the hair reflects to some extent the nutritional status and general health of the body. Thin, sparse hair and the loss of hair indicates a poor nutritional status. The deficiencies in diet, physical and mental illness of various kinds may also leads to premature graying of hair.

EN

- The hair follicles from which the hair grows produce oil which keeps the hair smooth. The sweat and the dead skin cells come off the scalp.
- The oil, sweat and dead cells all add together and can make the hair greasy and look dirty unless it is washed regularly.

S.No	Name of	the	Causative	Impacts/	Remedial
	Disease		Agents	Consequences	measures
1	Night Blindness		A lack o	f Makes it hard	Eat food rich
	_		vitamin A,	a to see well at	in anti –
			disorder c	f night or in	oxidant,
			the cells i	n poor light.	vitamins

				CHENNAI
		your retina		and
				minerals.
2	Conjunctivitis	Caused by a	One or both	Antibiotic
	(Pink eye)	virus and	eyes can be	eye drops or
		bacteria	affected.	ointments,
			Highly	home
			contagious;	remedy.
			can be spread	
			by	
			contamination	
			and sneezing	
3	Color blindness	Genetic	Difficulty	There is no
		condition	distinguishing	known cure
			between	for colour
			colours	blindness.
			Inability to see	contact
			shades or	lenses and
			tones of the	glasses with
	TA IN		same colour.	filters

To keep the hair clean and healthy:

- The regular hair wash and massage of the scalp will remove the dead skin cells, excess oil and dust.
- Rinsing the hair well with clear water and using good toothed comb for hair dressing is highly essential for their maintenance.

Diseases

- A disease is the functional or physical change from a normal state that affects the health of a person by causing disability or discomfort. The following are reasons that could leads to the development of disease in an individual.
- Infection caused by disease-causing microbes
- Lack of balanced diet
- > Poor lifestyle and unhealthy habits
- Malfunctioning of one or more body parts or organs.



Different kinds of sickness and their causes

• The prevention and treatment of sickness can be considered in two groups for their better understanding. They are, communicable and non-communicable disease.

Communicable Diseases

• Communicable diseases are those that spread from one person to another. Healthy persons must be protected from people with communicable diseases. Diseases spread through contaminated air, water, food or vectors (insects and other animals).

Diseases Caused by Bacteria

• Communicable disease (like tuberculosis, Cholera and Typhoid), which are caused by microbes and spread through air, water and some other organisms also.

NTRE

Tuberculosis:

• TB is caused by Mycobacterium tuberculae and spreads from one person to another person through air by spitting and prolonged contact with sharing materials of the patient. The symptoms are fever, weight loss, chronic cough, bloody spitting and difficulty in breathing.

Prevention and treatment

- ➢ BCG vaccination,
- Giving special attention to the patient,
- Regular medication like DOT

Cholera:

• Cholera is caused by Vibrio cholera and spread through the consumption contaminated food or water. The symptoms of Cholera is Vomiting, severe diarrhea and cramps in legs



Prevention and treatment

- Good hygienic practices like, washing hands before eating.
- Avoid eating uncovered food from street vendoers.
- Drinking boiled water.
- Getting Vaccination against cholera

Typhoid:

• Typhoid is caused by Salmonella typhi and spreads by contaminated food and water. The symptoms are Anorexia, headache, rashes on abdomen, dysentery and high fever up to 1040F.

Prevention and treatment

- Drinking boiled clean water
- Proper disposal of sewage
- ➢ Vaccination

Disease Caused by Virus

• Viral diseases are extremely widespread infections caused by many type of viruses. In this lesson you will learn about some disease caused by viruses like, Hapatitis, Chickenpox and Rabies.

Hepatitis

• Hepatitis is one of the most dangerous and fatal diseases caused by Hepatitis virus- A, B, C, D, E. Its mode of transmission is Contaminated water, sharing of needles and blood transfusion. The symptoms of hepatitis is loss of appetite, (Anorexia), vomiting, eyes and urine in yellow color.

Prevention and treatment

- Drinking boiled water,
- Proper cleaning of hands



Chickenpox

• Chickenpox (chicken pox), also known as varicella, is a highly contagious infection caused by the varicella zoster virus. This disease spreads through air and contact with an infected person. Its symptoms is appearance of rashes on the whole body, fever, headache and tiredness.

Prevention and treatment

- The chickenpox (varicella) vaccine is the best way to prevent chickenpox
- Special attention should be given to the infected persons

Rabies

Rabies is a fatal disease. Which is transmitted by the bite of the infected dog, rabbit, monkey, cat etc. The virus present in the saliva of dog enters the brain via neurons. The symptoms of rabies is hydrophobia (extreme fear for water), fever for 2 – 12 weeks and exaggerations in behavior.

Prevention and treatment

- In early stages rabies is very difficult to detect
- After an animal is bitten it usually takes two to twelve weeks to shows any symptoms and it may take as long as two years also.
- Fatality can be prevented by timely vaccination before the onset of symptoms.

Vaccine

A vaccine is a biological preparation that provides active acquired immunity to a particular disease. Vaccines like (BCG, Polio, MMR) are given at early child wood to protect from other diseases

Non-Communicable diseases

• Communicable diseases do not spread from person to person. They are caused by other factors. Therefore, it is important to know which sickness are communicable and which are not. They are never caused



by germs, bacteria, or other living organisms that infect the body. Antibiotics, or medicines that fight against germs do not help to cure non-communicable diseases.

Problems caused by wearing out of body parts:

Rheumatism, heart attack, epileptic seizures, stroke, migraine headach. Cataract and cancer

Problems caused by external harmful agents entering the body:

Allergies, asthma, poisons, snakebite, cough from smoking, stomach ulcer, alcoholism.

Problems caused by a lack of trace elements in the body:

Anemia, pellagra, night blindness and xerophthalmia, goiter and hypothyroidism.

Problems caused by Malnutrition.

Nutritious food is needed for a person to grow well, work hard, and stay healthy. Many common sicknesses are caused by malnutrition.

Leucoderma is a non – communicable diseases caused by partial or total loss of pigmentation in the skin. (melanin pigment). This condition affects any age, gender and ethnicity. There is no cure. It does not spread by touching, sharing food and sitting together.

Specific health problems of children

Anaemia

• It is caused by eating food with less iron content and can also caused due to feeding some other foods instead of breast milk. Severe anemia in children may leads to hookworm infection, chronic diarrhoea and dysentery. In recent day school going children, especially the girls are affected by anemia. The Government of Tamil Nadu provides weekly iron folic tablets to all the girls in the schools of all areas.



The signs of anemia are:

- Pale or transparent skin, The inner surface of eye lids are pale, white fingernails, pale gums, weakness and fatigue.
- In severe cases, face and feet may be swollen, the heartbeat is rapid and with shortness of breath.
- Children and women who eats mud are usually anemic.

Consuming iron containing food Sources

- Food Moringa leaves, Dates, Liver (Sheep and Chicken), Green, green leafy vegetables like beans, peas, lentils and Greed banana.
- > Pills Cod liver oil tablet, Ferrous sulfate

As a general rule, iron supplements should be given orally, not to be injected, because it leads to dangerous

Safety and First Aid

• First aid is the immediate treatment given to the victim of trauma or sudden illness before medical help is made available.

The first aid is

- ➢ To save the life
- To prevent further bleeding and determine the condition of the patient
- ➢ To relieve the pain
- > To provide a medical care available at the earliest

Burns

• The tissue damage caused by heat, chemical, electricity, sunlight or nuclear radiation is known as Burns. Mostly burns are caused by scalds, building fires, flammable liquid and gases. There are three types of burns, according to degree of burning.



- First-degree burns affect only the outer layer (called the epidermis) of the skin
- Second-degree burns damage the epidermis and the layer beneath it (called the dermis)
- Third-degree burns involve damage or complete destruction of the skin to its full depth and damage to underlying tissues also. People who experience such burns often require skin grafting.

First Aid for Burning

• In case of minor burns, the affected area should be washed with cold water and an antiseptic cream should be applied. In case of severe burns, where deeper layers of tissues get destroyed and blisters appear, use of water should be avoided. The burnt area should be covered with a clean non- sticking cloth or bandages. Larger burns need immediate medical attention.

Cut and Scratches

- Cuts and scratches are areas of damage on the surface of the skin. A cut is a line of damage that can go through the skin and into the muscle tissues below, whereas a scratch is surface damage that does not penetrate the lower tissues.
- Cuts and scratches may bleed or turn red, become infected and leave scars.

First aid for cuts

• For minor cuts, the affected area should be washed with cold running water and cleaned with an antiseptic liquid. Then apply an antiseptic cream on the wound and sterilized bandage should be tied to prevent infection. If the cut is deep, a clean cotton pad should be placed on the cut and pressed, and the injured person should be taken to a doctor immediately.



Basic cleanliness and protection.

- The most important thing is to help anybody, but you must also protect yourself from HIV and other blood-borne diseases when you help someone who is bleeding. You should wear gloves or a clean plastic bag on your hands.
- Be careful not to prick yourself with needles or other sharp objects around the person you are helping.





9th Full Book

Unit 21- Nutrition & Health

Introduction

- Food is the basic necessity of life. Food is defined as any substance of either plant or animal origin consumed to provide nutritional support for an organism. It contains essential nutrients that provide energy, helps in normal growth and development, repair the worn out tissues and protect the body from diseases. Food contamination with microorganisms is a major source of illness either in the form of infections or poisoning. Food safety is becoming a major concern these days.
- Adulteration of foodstuffs is commonly practiced in India by traders. Food is contaminated or adulterated from production to consumption for financial gain. The physiological functions of a consumer are affected due to either addition of a deleterious substance or the removal of a vital component. Food laws have come into existence to maintain the quality of food produced in our country.

Classes of Nutrients

Nutrients are classified into the following major groups as given below.

- Carbohydrates
- > Proteins
- ➤ Fats
- ➢ Vitamins
- ➤ Minerals



Carbohydrates

- Carbohydrates are organic compounds composed of carbon, hydrogen and oxygen. Carbohydrate is an essential nutrient which provides the chief source of energy to the body. Glucose, sucrose, lactose, starch, cellulose are examples for carbohydrates.
- Carbohydrates are classified as monosaccharide (Glucose), disaccharide (Sucrose) and polysaccharide (Cellulose). The classification is based on the number of sugar molecules present in each group.

Proteins

- Proteins are the essential nutrients and also the building blocks of the body. They are essential for growth and repair of body cells and tissues. Proteins are made of amino acids.
- Essential amino acids are those that cannot be biosynthesized by the body and must be obtained from the diet. The nine essential amino acids are phenylalanine, valine, threonine, tryptophan, methionine, leucine, isoleucine, lysine and histidine.

Fats

- Fat in the diet provides energy. They maintain cell structures and are involved in metabolic functions.
- Essential fatty acids cannot be synthesized in the body and are provided through diet. Essential fatty acids required in human nutrition are omega fatty acids.

Vitamins

• Vitamins are the vital nutrients, required in minute quantities to perform specific physiological and biochemical functions.

Human skin can synthesize Vitamin D when exposed to sunlight (especially early morning). When the sun rays falls on the skin dehydro



cholesterol is converted into Vitamin D. Hence, Vitamin D is called as Sunshine vitamin. Vitamin D improves bone strength by helping body to absorb calcium.

Minerals

- Minerals are inorganic substances required as an essential nutrient by organisms to perform various biological functions necessary for life. They are the constituents of teeth, bones, tissues, blood, muscle and nerve cells.
- The macrominerals required by the human body are calcium, phosphorus, potassium, sodium and magnesium. The microminerals required by the human body also called trace elements are sulfur, iron, chlorine, cobalt, copper, zinc, manganese, molybdenum, iodine and selenium.

Major food Stuffs	Dietary sources	Daily requirements
	V L	(grams)
Carbohydrates	Honey, Sugarcane,	150-200
TI	fruits, whole grains,	
2	starchy vegetables,	
	rice	
Proteins	Legumes, pulses ,	40
	nuts, soya bean, green	
	leafy vegetables, fish,	
	poultry products, egg,	
	milk and dairy	
	products	
Fats	Egg Yolk, saturated	35
	oil, meat	

Dietary sources of major foodstuffs

Vitamins-Dietary sources, Deficiency disorders and Symptoms

VitaminsDietary sourcesDeficiency disordersSymptoms					
Fat Soluble Vitamins					

A(Retinol)Papaya, leafy vegetables, fish liver oil, egg yolk, liver, dairy productsNyctalopia (NightCornea unabl to see in th night (dir light) scal skin.Vitamin D (Calciferol)Egg, liver, dairy products, Fish, Synthesized by the skin in sunlightRickets children)Image: Cornea unabl to see in th night (dir light)Vitamin (Tocopherol)E whole synthesized bi the skin in sunlightSterility in rats, Reproductive abnormalitiesSterility ir rats, Reproductive abnormalitiesVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is prevented bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.				CHENNAL
vegetables, fish liver oil, egg yolk, liver, dairy products(Night blindness)to see in th night (dir light) scal skin.Vitamin D (Calciferol)Egg, dairy products, Fish, Synthesized by the skin in sunlightRickets children)(in Bow defective rib development of pigeon chest.Vitamin (Tocopherol)E whole synthesized bi wegetable oil, meat, wegetables, Quinone)Sterility in rats, Reproductive abnormalitiesSterility ir rats, Reproductive abnormalitiesVitamin K (Derivative Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is prevented bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberi (Cheilosis)Degenerative eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	Vitamin	Carrot,	Xerophthalmia	Dryness of
fish liver oil, egg yolk, liver, dairy productsblindness)night (dir light) scal skin.Vitamin D (Calciferol)Egg, liver, dairy products, Fish, Synthesized by the skin in sunlightRickets (in children)Bow legg defective ribb development of pigeon chest.Vitamin (Tocopherol)E Whole Wheat, meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalitiesSterilityVitamin K (Derivative of Quinone)Leafy vegetables, milkBlood Clotting is bleeding due t delayed bloo clotting.Excessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberi changes in th nerves, muscle become weak Paralysis.Degenerative changes in th nerves, firstion i inflammation of lips, fissure in the corner of the mouth.	A(Retinol)	Papaya, leafy	Nyctalopia	Cornea unable
egg yolk, liver, dairy productslight)scal skin.Vitamin D (Calciferol)Egg, liver, dairy products, Fish, Synthesized by the skin in sunlightRickets (in children)Bow legg defective rib development of pigeon chest.Vitamin (Tocopherol)E Whole Wheat, meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalitiesSterility in rats, Reproductive abnormalitiesVitamin K (Derivative of Quinone)Leafy Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.		vegetables,	(Night	to see in the
egg yolk, liver, dairy productslight) skin.scal skin.Vitamin D (Calciferol)Egg, dairy products, Fish, Synthesized by the skin in sunlightRickets children)Bow defective ribb development of pigeon chest.Vitamin (Tocopherol)E whole Wheat, meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalitiesSterilityVitamin K (Derivative of Quinone)Leafy Soyabeans, milkBlood Clotting is bleeding due t clotting.Excessive bleeding due t clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis inflammation of lips, fissure in the corner of the mouth.		fish liver oil,	blindness)	night (dim
dairy productsskin.Vitamin D (Calciferol)Egg, liver, dairy products, Fish, Synthesized by the skin in sunlightRickets (in children)Bow legg defective rib development of pigeon chest.Vitamin (Tocopherol)E meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalities preventedSterilityVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.		egg yolk, liver,	,	U
(Calciferol)dairy products, Fish, Synthesized by the skin in sunlightchildren)defective ribs development of pigeon chest.Vitamin (Tocopherol)EWhole Wheat, meat, wegetable oil, milk.Sterility in rats, abnormalitiesSterilityVitamin K (Derivative Quinone)Leafy vegetables, milkBlood Clotting is preventedExcessive bleeding due t delayed blood clotting.Vitamin B1 (Thiamine)Whole grains, Peast, eggs, liver, sprouted pulsesBeriberiDegenerative eyes, dry skir inflammation of lips, fissure inflammation of lips, fissure in the corner of the mouth.		dairy products		skin.
products, Fish, Synthesized by the skin in sunlightdevelopment of pigeon chest.Vitamin (Tocopherol)EWhole Wheat, meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalitiesSterilityVitamin K (Derivative Quinone)Leafy of Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of lips, fissure in the corner of the mouth.	Vitamin D	Egg, liver,	Rickets (in	Bow legs,
Synthesized by the skin in sunlightpigeon chest.Vitamin (Tocopherol)EWhole Wheat, meat, meat, milk.Sterility in rats, Reproductive abnormalities preventedSterilityVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed blood clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of lips, fissure in the cornerVitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.	(Calciferol)	dairy	children)	defective ribs,
the skin in sunlightthe skin in sunlightVitamin (Tocopherol)E whole Wheat, meat, wegetable oil, milk.Sterility in rats, Reproductive abnormalitiesSterilityVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.		products, Fish,		development of
SunlightSunlightVitamin (Tocopherol)EWhole Wheat, meat, wegetable oil, abnormalities milk.Sterility in rats, Reproductive abnormalities prevented bleeding due t delayed bloo clotting.Vitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is prevented clotting.Excessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.		Synthesized by		pigeon chest.
Vitamin (Tocopherol)EWhole Wheat, meat, wegetable milk.Sterility in rats, Reproductive abnormalitiesSterility mats, Reproductive abnormalitiesVitamin K (Derivative Quinone)Leafy vegetables, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Water Soluble VitaminsDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, liver, green vegetables, whole grains liver, green vegetables, whole grainsAriboflavinosis inflammation of lips, fissure in the corner of the mouth.		the skin in		
(Tocopherol)meat, vegetable oil, milk.Reproductive abnormalities milticsVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed blood clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis inflammation of lips, fissure in the corner of the mouth.		sunlight		
(Tocopherol)meat, vegetable oil, milk.Reproductive abnormalitiesVitamin K (Derivative of Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed bloo clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weal Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.	Vitamin E	Whole Wheat,	Sterility in rats,	Sterility
milk.LeafyBlood Clotting isExcessiveVitamin K (Derivative Quinone)LeafyBlood Clotting isExcessiveSoyabeans, milkpreventedbleeding due tQuinone)Soyabeans, milkdelayed blood clotting.Water Soluble VitaminsVitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	(Tocopherol)	meat,	Reproductive	-
Vitamin K (Derivative Quinone)Leafy vegetables, Soyabeans, milkBlood Clotting is preventedExcessive bleeding due t delayed blood clotting.Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis of lips, fissure in the corner of the mouth.		vegetable oil,	abnormalities	
(Derivative Quinone)of vegetables, Soyabeans, milkprevented preventedbleeding due t delayedWater Soluble VitaminsVitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.		U		
Quinone)Soyabeans, milkdelayed blood clotting.Water Soluble VitaminsVitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	Vitamin K	Leafy	Blood Clotting is	Excessive
milkclotting.Water Soluble VitaminsVitamin B1Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	(Derivative of	vegetables,	prevented	bleeding due to
Water Soluble VitaminsVitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	Quinone)	Soyabeans,		delayed blood
Vitamin B1 (Thiamine)Whole grains, Yeast, eggs, liver, sprouted pulsesBeriberiDegenerative changes in th nerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.		milk	TI	clotting.
(Thiamine)Yeast, eggs, liver, sprouted pulseschanges in th nerves, muscle become weak Paralysis.Vitamin B2Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.		Water Solubl	le Vitamins	
liver, sprouted pulsesnerves, muscle become weak Paralysis.Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation i eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	Vitamin B1	Whole grains,	Beriberi	Degenerative
pulsesbecome weak Paralysis.Vitamin B2Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation is eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	(Thiamine)	Yeast, eggs,		changes in the
Vitamin B2Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	c T	liver, sprouted		nerves, muscles
Vitamin B2 (Riboflavin)Milk, eggs, liver, green vegetables, whole grainsAriboflavinosis (Cheilosis)Irritation eyes, dry skir inflammation of lips, fissure in the corner of the mouth.	D	pulses		become weak,
(Riboflavin)liver, green vegetables, whole grains(Cheilosis)eyes, dry skir inflammation of lips, fissure in the corner of the mouth.				Paralysis.
vegetables, whole grains in the corner of the mouth.	Vitamin B2	Milk, eggs,	Ariboflavinosis	Irritation in
whole grains of lips, fissure in the corner of the mouth.	(Riboflavin)	liver, green	(Cheilosis)	eyes, dry skin,
in the corner of the mouth.		vegetables,		inflammation
of the mouth.		whole grains		of lips, fissures
				in the corners
				of the mouth.
VITAMIN B3 MIIK, eggs, Pellagra Inflammation	Vitamin B3	Milk, eggs,	Pellagra	Inflammation
(Niacin) liver, lean of skin, loss of	(Niacin)	liver, lean		of skin, loss of
meat, ground memory,		meat, ground		
nuts, bran diarrhoea		nuts, bran		diarrhoea
Vitamin B6Meat,fish,DermatitisScalyskir	Vitamin B6	Meat, fish,	Dermatitis	Scaly skin,
(Pyridoxine) eggs, germs of nervous	(Pyridoxine)	eggs, germs of		nervous
grains and disorders.				disorders.



	cereals, rice polishings.		
Vitamin B12	Milk, meat,	Pernicious	Decrease in red
(Cyanocobalamine)	liver, pulses,	anaemia	blood cell
	cereals, fish		production,
			degeneration of
			spinal cord.
Vitamin C	Leafy	Scurvy	Swollen and
(Ascorbic acid)	vegetables,		bleeding gums,
	sprouts, citrus		delay in
	fruits like		healing of
	gooseberry		wounds, teeth
	(Amala),		and bones
	lemon, orange		malformed.

Minerals - Dietary sources, Functions and Deficiency disorders

Minerals	300008		Deticionau
	Sources	Functions	Deficiency
			disorders
	Macro	nutrients	
Calcium	Dairy Products,	Constituent of	Bone
	beans, cabbage,	bones and enamel	deformities,
	eggs, fish	of teeth, clotting	poor skeletal
S		of blood and	growth,
		controls muscle	Osteoporosis in
		contraction.	adults.
Sodium	Common Salt	Maintains fluid	Muscular
		balance and	cramps, nerve
		involved in	impulses do not
		neurotransmission	get transmitted.
Potassium	Banana, Sweet	Regulates nerve	Muscular
	Potato, nuts,	and muscle	fatigue, nerve
	whole grains,	activity	impulses do not
	citrus fruits		get transmitted.
Micro nutrients			
Iron	Spinach, dates,	Important	Anaemia
	greens,	component of	
	broccoli, whole	haemoglobin	
	cereals, nuts,	-	



	fish, liver				
Iodine	Milk,	Seafood,	Formation	of	Goitre
	Iodised salt		thyroid hormones.		

Protein Energy Malnutrition (PEM)

- Absence of certain nutrients in our daily diet over a long period of time leads to deficiency diseases. This condition is referred as Malnutrition. Deficiency of proteins and energy leads to severe conditions like: Kwashiorkar and Marasmus.
- Kwashiorkar: It is a condition of severe protein deficiency. It affects children between 1-5 years of age, whose diet mainly consists of carbohydrates but lack in proteins.
- Marasmus: It usually affects infants below the age of one year when the diet is poor in carbohydrates, fats and proteins.

Food Hygiene

- Poor personal hygiene may allow pathogenic microorganisms to cause food spoilage Food spoilage is an undesirable change in the normal state of food and is not suitable for human consumption. Signs of food spoilage include a changes in appearance, colour, texture, odour and taste. Factors responsible for Food Spoilage are given below.
- Internal **factors:** It include enzymatic activities and moisture content of the food.
- **External factors:** It include adulterants in food, contaminated utensils and equipment, unhygienic cooking area and lack of storage facilities.

Food Preservation

• Food preservation is the process of prevention of food from decay or spoilage, by storing in a condition fit for future use. Food is preserved to:



- increase the shelf life of food
- > retain the colour, texture, flavour and nutritive value
- ➢ increase food supply
- decrease wastage of food

Methods of Food Preservation

The various method of food preservation are explained below

- Drying: Drying is the process of preservation of food by removal of water/moisture content in the food. It can be done either by sundrying, (e.g. cereals, fish) or vacuum drying (e.g. milk powder, cheese powder) or hot air drying (e.g. grapes, dry fruits, potato flakes). Drying inhibits the growth of microorganism such as bacteria, yeasts and moulds.
- Smoking: In this process, food products like meat and fish are exposed to smoke. The drying action of the smoke tends to preserve the food.
- Irradiation: Food irradiation is the process of exposing food to optimum levels of ionizing radiations like x-rays, gamma rays or UV rays to kill harmful bacteria and pests and to preserve its freshness.
- Cold **storage:** It is a process of storing the perishable foods such as vegetables, fruits and fruit products, milk and milk products etc. at low temperature. Preserving the food products at low temperature slows down the biological and chemical reactions and prevents its spoilage.
- Freezing: Freezing is one of the widely used methods of food preservation. This process involves storing the food below 00C at which microorganisms cannot grow, chemical reactions are reduced and metabolic reactions are also delayed.

Pasteurization: Pasteurization is a process of heat treatment of liquid food products. e.g. For preservation of milk and beverages. This process also involves boiling of milk to a temperature of 63°C for about 30



minutes and suddenly cooling to destroy the microbes present in the milk.

Bananas are best stored at room temperature. When it is kept in a refrigerator, the enzyme responsible for ripening becomes inactive. In addition, the enzyme responsible for browning and cell damage becomes more active thereby causing the skin colour change from yellow to dark brown

• Canning: In this method of food preservation, most vegetables, fruits, meat and dairy products, fruit juices and some ready-to-eat foods are processed and stored in a clean, steamed air tight containers under pressure and then sealed. It is then subjected to high temperature and cooled to destroy all microbes.

Addition of Preservatives

• Food can be preserved by adding natural and synthetic preservatives.

TRE

Natural preservatives

- Some naturally available materials like salt, sugar and oil are used as food preservatives.
- Addition **of salt:** It is one of the oldest methods of preserving food. Addition of salt removes the moisture content in the food by the process of osmosis. This prevents the growth of bacteria and reduces the activity of microbial enzymes. Meat, fish, gooseberry, lemon and raw mangoes are preserved by salting. Salt is also used as a preservative in pickles, canned foods etc.
- Addition of sugar: Sugar/Honey is added as a preservative to increase the shelf life of fruits and fruit products like jams, jellies, squash, etc. The hygroscopic nature of sugar/honey helps in reducing the water content of food and also minimizing the process of oxidation in fruits.
- Addition of oil: Addition of oil in pickles prevents the contact of air with food. Hence microorganisms cannot grow and spoil the food.



Synthetic preservatives

• Synthetic food preservatives like sodium benzoate, citric acid, vinegar, sodium meta bisulphate and potassium bisulphate are added to food products like sauces, jams, jellies, packed foods and ready- to-eat foods. These preservatives delay the microbial growth and keep the food safe for long duration.

Food Adulteration

- Adulteration is defined as the addition or subtraction of any substance to or from food, so that the natural composition and the quality of food substance is affected. Adulterant is any material which is used for the purpose of adulteration.
- Some of the common adulterated foods are milk and milk products, cereals, pulses, coffee powder, tea powder, turmeric powder, saffron, confectionary, non-alcoholic beverages, spices, edible oils, meat, poultry products etc. The adulterants in food can be classified in three categories:
- Natural adulterants
- Incidental/unintentionally added adulterants
- Intentionally added adulterants.

Natural adulterants

• Natural adulterants are those chemicals or organic compounds that are naturally present in food. e.g. toxic substances in certain poisonous mushrooms, Prussic acid in seeds of apples and cherry, marine toxins, fish oil poisoning, environmental contaminants.

Incidental/unintentionally added adulterants

- These types of adulterants are added unknowingly due to ignorance or carelessness during food handling and packaging. It includes:
- Pesticide residues
- Droppings of rodents, insects, rodent bites and larva in food during its storage



Microbial contamination due to the presence of pathogens like Escherichia coli, Salmonella in fruits, vegetables, ready-to-eat meat and poultry products.

Intentionally added adulterants

- These adulterants are added intentionally for financial gain and have serious impact on the health of the consumers. These types of adulterants include:
- Additives and preservatives like vinegar, citric acid, sodium bicarbonate (baking soda), hydrogen peroxide in milk, modified food starch, food flavours, synthetic preservatives and artificial sweeteners.
- > Chemicals like calcium carbide to ripen bananas and mangoes.
- Non certified food colours containing chemicals like metallic lead are used to give colours to vegetables like green leafy vegetables, bitter gourd, green peas etc. These colours are added to give a fresh look to the vegetables.
- Edible synthetic wax like shellac or carnauba wax is coated on fruits like apple, pear to give a shining appearance.

Health Effects of Adulterated Foods

• Consumption of these adulterated foods may lead to serious health effects like fever, diarrhoea, nausea, vomiting, gastrointestinal disorders, asthma, allergy, neurological disorder, skin allergies, immune suppression, kidney and liver failure, colon cancer and even birth defects.

Food Quality Control

• The government always ensures that pure and safe food is made available to the consumers. In 1954, the Indian Government enacted the Food Law known as Prevention of Food Adulteration Act and the Prevention of Food Adulteration Rules in 1955 with the objective of



ensuring pure and wholesome food to the consumers and protect them from fraudulent trade practices.

• Minimum standards of quality for food and strict hygienic conditions for its sale are clearly outlined in the Act

A slogan From farm to plate, make food safe was raised on World Health Day (7th April 2015) to promote and improve food safety.

Food Quality Control Agencies

- ISI, AGMARK, FPO, FCI and other health departments enforce minimum standards for the consumer products. FCI (Food Corporation of India) was set up in the year 1965 with the following objectives:
- Effective price support operations for safeguarding the interest of farmers.
- Distributing food grains throughout the country.
- Maintaining satisfactory levels of operational and buffer stock of food grains to ensure national security.
- Regulate the market price to provide food grains to consumers at reliable price.



10th Full Book

Unit 21 - Health & Disease

Introduction

Abuses occur in a variety of forms and are deeply rooted in cultural, social and economic practices. Solving this global problem however requires a much better understanding of its occurrence, causes and consequences with context to sexual and childhood abuse, this is followed by substance abuse. Are people leading healthier lives in today's modern world than their generations did in the past? For instance, smoking cigarettes, alcohol addiction, use of drugs, eating high fat and cholesterol rich diets, excessive intake of junk foods, reduced physical activity are some of the risk factors for illness and early death. The role of behaviour in health has been receiving increased attention in countries around the world. The health habits of the individuals and their behaviour influence the development of chronic and fatal diseases such as diabetes, obesity, heart disease, cancer and AIDS. These conditions can be substantially reduced by adopting lifestyles that promote wellness and protect their health by taking nutritious diet, regular exercise and by avoiding drugs, alcohol and smoking.

Abuse and Types of Abuse

- Abuse refers to cruel, violent, harmful or injurious treatment of another human being. It includes physical, emotional or psychological, verbal, child and sexual abuses. Abuse can occur within the family and with people who are not associated with the family.
- These days the use of drugs, alcohol and tobacco has been increasing especially among teenagers and adolescents for adventure, excitement, curiosity and experimentation



Child Abuse

• Child abuse constitutes all forms of physical or emotional ill treatment, sexual abuse, exploitation resulting in child's ill health, survival and development. Physical abuse of a child is defined as those acts that cause physical harm such as threatening, beating, kicking and hitting the child.

Sexual Abuse

• Sexual harassment is a form of power and dominance of one person over another, which can result in harmful consequence to the victim. It refers to inappropriate or forced sexual contact. Adolescent girls and women encounter sexual harassment in different forms. Sexual abuse is more common at work places. Verbal remarks, comments, gestures and looks are the most common forms of abuse. This results in psychological distress, physical illness and eating disorders in the ENTRE affected individuals.

Child Sexual Abuse

- Children are considered soft targets for sexual abuse because they may not realize that they are being abused. Commonly, abusers are persons well known to the child, may even be living in the same locality. Abusers also bribe (use chocolates and toys) to lure children and take advantage of the child's innocence.
- Sexually abused children show symptoms of genital injury, abdominal pain, frequent urinary infection and behavioural problems.

Approaches for Protection of an Abused Child

- Measures adopted for monitoring and assessment of abused child who have undergone signs and symptoms of distress are:
 - ✓ **Child Helpline:** The Child Helpline provides a social worker who can assist the child by providing food, shelter and protection.



- ✓ Counselling the child: Psychologists and social workers should provide guidance, counselling and continous support to a victimized child.
- ✓ Family support: The victimized child should be supported by the family members. They should be provided with proper care and attention to overcome their sufferings.
- ✓ Medical care: A child victim of sexual offences should receive medical care and treatment from health care professionals to overcome mental stress and depression.
- ✓ Legal Counsel: The family or the guardian of the child victim shall be entitled to free assistance of a legal counsel for such offence.
- ✓ Rehabilitation: Enrolling in schools and resuming their education is an important step towards rehabilitation of the child. It is essential that the child's life is gradually returned to normal after the incidence of abuse.
- ✓ Community based efforts: Conducting awareness campaign on child abuse and its prevention.

Prevention of child sexual abuse

- The most important social policy proclaimed universally is the prevention of child abuse. Taking steps to prevent childhood sexual abuse is parental and institutional responsibility. Instructions to be given by parents and teachers to the child are.
- Do not talk to any suspected person or strangers and to maintain a distance.
- Not to be alone with unknown person.
- > To be careful while travelling alone in public or private transport.
- Not to receive money, toys, gift s or chocolates from known or unknown person to them without the knowledge of their parents.



- ▶ Not to allow known or unknown person to touch them.
- It is the responsibility of every individual living in a society to ensure a safe and protected environment for our children to enable them to live with dignity and free from any form of violence.

Drug, Alcohol and Tobacco Abuse

• The physical and mental dependency on alcohol, smoking and drugs is called addiction. The addictive potential of these substances pulls an individual into a vicious cycle leading to regular abuse and dependency. This is of serious concern because abuse of tobacco, alcohol or drugs produce many harmful effects in an individual, to the family and even to the society. This dangerous behavior pattern among youth can be prevented through proper guidance.

Drug Abuse

- Drugs are normally used for the treatment of disease on advice of a physician and withdrawn after recovery. A person who is habituated to a drug due to its prolonged use is called drug addict. This is called drug addiction or drug abuse.
- A drug that modifies the physical, biological, psychological or social behaviour of a person by stimulating, depressing or disturbing the functions of the body and the mind is called addictive drug. These drugs interact with the central nervous system and affect the individual physically and mentally.

Types of Drugs

• There are certain drugs called psychotropic drugs which acts on the brain and alter the behaviour, consciousness, power of thinking and perception. They are referred as mood altering drugs.

Drug Dependence

• Persons who consume these drugs become fully dependent on them, they cannot live without drugs. This condition is referred as drug dependence.



- > Physical and mental dependence
- Dependence on the drug for normal condition of well being and to maintain physiological state.
- Psychological dependence is a feel that drugs help them to reduce stress.
- International Day against Drug Abuse and Illicit Trafficking June 26.
- Narcotic Drugs and Psychotropic Substances Act was introduced in 1985.

Behavioural Changes of Drug Users

- Adverse effects of drug use among adolescents are
- > Drop in academic performance, absence from school or college.
- Lack of interest in personal hygiene, isolation, depression, fatigue and aggressive behaviour.
- > Deteriorating relationship with family and friends.
- Change in food and sleeping habits
- Fluctuation in body weight and appetite
- Always looking out for an easy way to get money for obtaining drugs.
- > Prone to infections like AIDS and Hepatitis-B.

World Health Organization (WHO) 1984 suggested the use of the term drug dependence in place of drug addiction or drug abuse

Drug De-addiction

• Management of de-addiction is a complicated and difficult task. The path to recovery of drug addicts is long and often slow.



- Family members, friends and society on the whole have a very important role to play.
 - ✓ Detoxification: The first phase of treatment is detoxification. The drug is stopped gradually and the addict is helped to overcome the withdrawal symptoms. The addict undergoes severe physical and emotional disturbance. This is taken care by specific medication.
 - ✓ Psychotherapy: Individual and group counselling is given by psychologists and counsellors. The treatment includes efforts to reduce the addict's stress, taught new ways to solve everyday's problems, adequate diet, rest and relaxation.
 - Counselling to family members: Social workers counsell family members in order to change the attitude of rejection so that the addict is accepted by the family and the society.
 - ✓ Rehabilitation: They are given proper vocational training so that they can lead a healthy life and become useful members of the society.

Tobacco Abuse

• Tobacco is obtained from the tobacco plant Nicotiana tobaccum and Nicotiana rustica. The dried and cured leaves of its young branches make the commercial tobacco used worldwide. Addiction to tobacco is due to 'nicotine' an alkaloid present in it. Nicotine is a stimulant, highly harmful and poisonous substance.

Tobacco Use

• Tobacco is used for smoking, chewing and snuffing. Inhaling tobacco smoke from cigars, cigarettes, bidis, pipes, hukka is called smoking. Tobacco in powder form is chewed with pan. When powdered tobacco is taken through nose, it is called snuffing.



Smoking Hazards and Effects of Tobacco

- When smoke is inhaled, the chemicals get absorbed by the tissues and cause the following harmful effects
- Benzopyrene and polycyclic hydrocarbons present in tobacco smoke is carcinogenic causing lung cancer.
- Causes inflammation of throat and bronchi leading to conditions like bronchitis and pulmonary tuberculosis.
- Inflammation of lung alveoli, decrease surface area for gas exchange and cause emphysema.
- Carbon monoxide of tobacco smoke binds to haemoglobin of RBC and decreases its oxygen carrying capacity causing hypoxia in body tissues.
- Increased blood pressure caused by smoking leads to increased risk of heart disease.
- Causes increased gastric secretion which leads to gastric and duodenal ulcers.
- Tobacco chewing causes oral cancer (mouth cancer).

Prevention of Smoking

• Knowing the dangers of smoking and chewing tobacco adolescents and the old people need to avoid these habits. Proper counselling and medical assistance can help an addict to give up the habit of smoking.

Alcohol Abuse

• The consumption of alcohol is a social evil practiced by the wealthier and poorer sections of the society. The dependence of alcohol is called alcoholism and the addict is termed as alcoholic. It is called alcohol abuse. Drinking of alcohol impairs one's physical, physiological and psychological functions.



Harmful Effects of Alcohol to Health

- Prolonged use of alcohol depresses the nervous system, by acting as a sedative and analgesic substance. Some of the harmful effects are
- Nerve cell damage resulting in various mental and physical disturbances
- Lack of co-ordination of body organs
- Blurred or reduced vision, results in road accidents
- > Dilation of blood vessels which may affect functioning of the heart
- Liver damage resulting in fatty liver which leads to cirrhosis and formation of fibrous tissues
- Body loses its control and consciousness eventually leading to health complications and ultimately to death.

Rehabilitation Measures for Alcoholics

- ✓ Education and counselling: Education and proper counselling will help the alcoholics to overcome their problems and stress, to accept failures in their life.
- ✓ Physical activity: Individuals undergoing rehabilitation should be channelized into healthy activities like reading, music, sports, yoga and meditation.
- ✓ Seeking help from parents and peer groups: When a problematic situation occurs, the affected individuals should seek help and guidance from parents and peers. This would help them to share their feeling of anxiety, wrong doing and get rid of the habit.
- ✓ Medical assistance: Individual should seek help from psychologists and psychiatrists to get relieved from this condition and to lead a relaxed and peaceful life.



• Alcohol de-addiction and rehabilitation programmes are helpful to the individual so that they could get rid of the problem completely and can lead a normal and healthy life.

Diseases and Disorders due to Lifestyle Modifications

• Diseases are prevalent in our society due to our improper way of living, conditions of stress and strain. These diseases are non-communicable and affect the person who are suffering from particular symptoms. It is an impairment of the body tissue or organ, disturbances in metabolic function which require modification of an individual's normal life.

Diabetes Mellitus

Diabetes mellitus is a chronic metabolic disorder. In Greek (Diabetes
 – running through; mellitus- sweet). It is characterised by increased
 blood glucose level due to insufficient, deficient or failure of insulin
 secretion. This is the most common pancreatic endocrine disorder.
 The incidence of Type-1 and Type-2 diabetes is increasing worldwide.

Type-1 Insulin Dependent Diabetes Mellitus (IDDM)

• IDDM accounts for 10 to 20% of the known diabetics. The condition also occurs in children (juvenile onset diabetes) and young adults, the onset is usually sudden and can be life threatening. This is caused by the destruction of β -cells of the pancreas. It is characterized by abnormally elevated blood glucose levels (hyperglycemia) resulting from inadequate insulin secretion.

Causes: Genetic inheritance and environmental factors (infections due to virus, acute stress) are the cause for this condition.

Type-2 Non-Insulin Dependent Diabetes Mellitus (NIDDM)

• This is also called as adult onset diabetes and accounting for 80 to 90% of the diabetic population. It develops slowly, usually milder and more stable. Insulin production by the pancreas is normal but its action is impaired. The target cells do not respond to insulin. It does not allow the movement of glucose into cells.



Causes: The causes are multifactorial which include increasing age (affecting middle aged and older people), obesity, sedentary life style, overeating and physically inactive.

Symptoms: Diabetes mellitus is associated with several metabolic alterations. The most important symptoms are

- Increased blood glucose level (Hyperglycemia)
- Increased urine output (Polyuria) leading to dehydration
- Loss of water leads to thirst (Polydipsia) resulting in increased fluid intake
- Excessive glucose excreted in urine (Glycosuria)
- > Excess hunger (Polyphagia) due to loss of glucose in urine.
- Fatigue and loss of weight

According to WHO recommendation, if the fasting blood glucose is greater than 140 mg/dl or the random blood glucose is greater than 200 mg /ml on more than two occasions, diagnosis for confirming diabetes is essential.

Prevention and Control of Diabetes

• Diet, hypoglycemic drugs, insulin injection and exercise are the management options based on the type and severity of the condition. The overall goal of diabetes management is to maintain normal blood glucose level.

Differences bet	ween Type-1 a	nd Type-2 Diabete	s Mellitus
	21	21	

Factors	Dependent Diabetes	Type-2 Non-Insulin Dependent Diabetes Mellitus (NIDDM)
Prevalence	10-20%	80 - 90%
Age of onset	Juvenile onset (< 20 Years)	Maturity onset (



		CHENNAI
		>30 Years)
Body Weight	Normal or	obese
	underweight	
Defect	Insulin deficiency due	Target cells do
	to destruction of β –	respond to insulin
	cells.	
Treatment	Insulin administration	Can be controlled by
	is necessary	diet, exercise and
	-	medicine

- ✓ Dietary management: Low carbohydrate and fibre rich diets are more appropriate. Carbohydrates should be taken in the form of starch and complex sugars. Refined sugars (sucrose and glucose) should be avoided. Diet comprising whole grains, millets (jowar, bajra, ragi), green leafy vegetables, wheat and unpolished rice should be included in diet regularly. Carbohydrates is maintained to about 50- 55% of the total calories. High protein content of 10-15% of the total intake is required to supply essential amino acids. Fat content in the diet should be 15-25% of the total calories. Saturated fat intake should be reduced. Polyunsaturated fatty acid content should be higher.
- ✓ Management with insulin: Commercially available insulin preparations (short and long acting) are also used to maintain blood glucose levels.
- Physical activity: Exercise plays an important role in facilitating a good control of diabetes, in addition to strengthening and toning up the muscles.
- ✓ Education and Awareness: People with diabetics should be educated on the nature of disease they have and the possibility of complications of the disease, if blood sugar is not kept under control. Instructions regarding diet, exercise and drugs should be explained.



Obesity

- Obesity is the state in which there is an accumulation of excess body fat with an abnormal increase in body weight. Obesity is a complex multifactorial chronic disease developing from influence of social, behavioural, psychological, metabolic and cellular factors.
- Obesity occurs if intake of calories is more than the expenditure of energy. Over weight and obesity are conditions where the body weight is greater than the mean standard weight for age and height of an individual. Body mass index (BMI) is an estimate of body fat and health risk.

BMI = Weight (kg) / Height (m)^{2.}

Causes and risk factors: Obesity is due to genetic factors, physical inactivity, eating habits (overeating) and endocrine factors. Obesity is a positive risk factor in development of hypertension, diabetes, gall bladder disease, coronary heart disease and arthritis.

Prevention and Control of Obesity

- Diet Management: Low calorie, normal protein, vitamins and mineral, restricted carbohydrate and fat, high fiber diet can prevent overweight. Calorie restriction for weight reduction is safe and most effective.
- Physical exercise: A low calorie diet accompanied by moderate exercise will be effective in causing weight loss. Meditation, yoga and physical activity can also reduce stress related to overeating.

Heart Disease

- Cardiovascular disease (CVD) is associated with diseases of the heart and blood vessels. Coronary heart disease (CHD) is the most common form and is caused by deposition of cholesterol in the blood vessels.
- It usually develops slowly over many years beginning from childhood, they may form a fatty streak to a fibrous complicated plaque. It leads to the narrowing of blood vessels leading to



atherosclerosis in the large and medium sized arteries that supply the heart muscle with oxygen. It leads to sudden ischemia (deficient blood supply to heart muscle) and myocardial infarction (death of the heart muscle tissue).

- **Risk factors:** Hypercholesterolemia (High blood cholesterol) and high blood pressure (Hypertension) are the major causes and contributing factors for heart disease and if untreated may cause severe damage to brain, kidney and eventually lead to stroke.
- **Causes:** Heredity (family history), diet rich in saturated fat and cholesterol, obesity, increasing age, cigarette smoking, emotional stress, sedentary lifestyle, excessive alcohol consumption and physical inactivity are some of the causes.
- **Symptoms:** Shortness of breath, headache, tiredness, dizziness, chest pain, swelling of leg, and gastrointestinal disturbances.

HDL (High Density Lipoprotein) or "good" cholesterol lowers risk of heart disease while LDL (Low Density Lipoprotein) or "bad" cholesterol increases risk of heart disease.

Prevention and Control of Heart Disease

- **Diet management:** Reduction in the intake of calories, low saturated fat and cholesterol rich food, low carbohydrates and common salt are some of the dietary modifications. Diet rich in polyunsaturated fatty acids (PUFA) is essential. Increase in the intake of fibre diet, fruits and vegetables, protein, minerals and vitamin are required.
- **Physical activity:** Regular exercise, walking and yoga are essential for body weight maintenance
- Addictive substance avoidance: Alcohol consumption and smoking are to be avoided.

Cancer

• Cancer causes about 4 million deaths annually throughout the world. In India more than one million people suffer from cancer. Cancer is



derived from Latin word meaning crab. The study of cancer is called Oncology (Oncos-Tumor).

- Cancer is an abnormal and uncontrolled division of cells that invade and destroy surrounding tissue forming a tumor or neoplasm (new growth). It is a heterogenous group of cells that do not respond to the normal cell division.
- The cancerous cells migrate to distant parts of the body and affect new tissues. This process is called metastasis. The frequent sites of metastasis are lungs, bones, liver, skin and brain.

Types of Cancers

- Cancers are classified on the basis of the tissues from which they are formed.
- Carcinomas arise from epithelial and glandular tissues. They include cancers of skin, lung, stomach and brain. About 85% of the tumours are carcinomas
- Sarcomas are occur in the connective and muscular tissue. They include the cancer of bones, cartilage, tendons, adipose tissue and muscles. These form 1% of all tumours.
- Leukaemia are characterized by an increase in the formation of white blood cells in the bone marrow and lymph nodes. Leukaemia are called blood cancers. Most common type of cancer which also affect children below 15 years of age.

Carcinogenic Agents

- Cancer causing agents are called carcinogens. They are physical, chemical agents, ionizing radiations and biological agents.
 - ✓ Physical Irritant: Heavy smoking causes lung cancer and cancers of oral cavity, pharynx (throat) and larynx. Betel and tobacco chewing causes oral cancer. Excessive exposure to sunlight may cause skin cancer.



- Chemical agents: Nicotine, caffeine, products of combustion of coal and oil, pesticides, asbestos, nickel, certain dyes and artificial sweetners induce cancer
- ✓ Radiations: Ionizing radiations like X-rays, gamma- rays, radioactive substances and non-ionising radiations like UV rays cause DNA damage leading to cancer.
- ✓ Biological agents: Cancer causing viruses are called oncogenic viruses.

Treatment of Cancer

The treatment of cancer involves the following methods:

- Surgery: Tumours are removed by surgery to prevent further spread of cancer cells.
- Radiation therapy: Tumour cells are irradiated by lethal doses of radiation while protecting the surrounding normal cells.
- Chemotherapy: It involves administration of anticancerous drugs which prevent cell division and are used to kill cancer cells.
- Immunotherapy: Biological response modifiers like interferons are used to activate the immune system and help in destroying the tumors.

Preventive measures for Cancer

- Cancer control programmes should focus on primary prevention and early detection.
- To prevent lung cancer tobacco smoking is to be avoided and protective measures to be taken against exposure to toxic pollutants of industries. Excessive exposure to radiation is to be avoided to prevent skin cancer.



AIDS (Acquired Immunedeficiency Syndrome)

• AIDS is a severe viral disease and caused by Human Immunodeficiency Virus (HIV). It is a condition in which immune system fails and suppress the body's disease fighting mechanism. They attack the lymphocytes and the affected individual is prone to infectious diseases.

Dr. Suniti Solomon, pioneered HIV research and treatment in India. She set up the first voluntary testing and counselling centre and an AIDS Research group in Chennai during 80's. Her team was the first to document evidence of HIV infection in India in 1985 (First Indian AIDS patient in Chennai).

Transmission of HIV

• AIDS virus has been found in urine, tears, saliva, breast milk and vaginal secretions. The virus is transmitted by an infected patient who comes in contact with blood of a healthy person. HIV/AIDS is not transmitted by touch or any physical contact. It spreads through contact of body fluids or blood.

HIV is transmitted generally by

- Sexual contact with infected person
- Use of contaminated needles or syringes especially in case of intravenous drug abusers
- > By transfusion of contaminated / infected blood or blood products
- > From infected mother to her child through placenta.

Symptoms and Treatment of AIDS

Symptoms: Infected individuals become immunodeficient. The person becomes more susceptible to viral, bacterial, protozoan and fungal infections. Swelling of lymph nodes, damage to brain, loss of memory, lack of appetite and weight loss, fever, chronic diarrhoea, cough, lethargy, pharyngitis, nausea and headache.

Diagnosis: The presence of HIV virus can be confirmed by Western Blot analysis or Enzyme Linked Immunosorbent Assay (ELISA)



Treatment: Anti-retroviral drugs and immunestimulative therapy can prolong the life of the infected person.

Prevention and Control of AIDS

- The following steps may help in controlling and prevent the spreading of HIV infection
- Screening of blood from blood banks for HIV before transfusion.
- Ensuring the use of disposable needles and syringes in hospitals and clinics.
- Advocating safe sex and advantages of using condoms.
- Creating awareness campaign and educating people on the consequences of AIDS.
- Persons with HIV/AIDS should not be isolated from the family and society.



11th Zoology

Chapter 5 – Digestion and Absorption

Nutrients, Vitamins and Minerals

- Food comprises of macronutrients and micronutrients. The nutrients required in larger quantities are called macronutrients, whereas those required in small quantities are called micronutrients. Essential nutrients cannot be synthesized by the body; they have to be included in the diet. Macronutrients are lipids, carbohydrates, proteins and the micronutrients are vitamins and minerals. Water plays an important role in the metabolic processes and prevents dehydration of the body.
- Intake of too much of food or lesser amount of food than the basic requirement is called malnutrition. A diet which can provide all the metabolic requirements of the body in a right proportion is called balanced diet. That means it should contain carbohydrates and fats for energy yielding, proteins for growth and replacement; and vitamins, minerals and water for physiological regulation.

Vitamins:

• Vitamins are naturally occurring organic substances regularly needed in minute quantities for maintaining normal health as metabolic regulators. The identified vitamins are classified as fat soluble (A,D,E and K) and vitamin B and vitamin C are water soluble. Vitamin A, D, E and K, if consumed beyond required level may cause defects, commonly referred to as hypervitaminosis.

Minerals:

• These are the inorganic chemical elements, i.e., Ca, Fe, I, K, Mg, Na, P, S, etc needed for regulation of various physiological functions. These can be classified into major minerals (Na, P, K, Ca, Mg, S, Cl) and others are trace minerals such as Fe, Cu, Zn, Co, Mn, I, and fluorine.



Sodium ions are more abundant than any other cation in the body fluids.

N.I. Lunin discovered vitamins but the name vitamin was given by Dr. Funk (1912). The first vitamin isolated was B1 by Dr.Funk. The first vitamin produced by fermentation process using, Acetobacter bacteria is Vitamin C.

Fat soluble vitamins		
Vitamins	Functions	Symptoms of
		Deficiency
A (Retinol)/	Plays a vital role in	Night blindness
Antixerophthalmic	visual perception.	(Nyctalopia),
vitamin	Maintenance and	Xerophthalmia
	growth of epithelial	(drying of eyeballs),
	tissue.	Bitot's spot in the
		cornea, Dermatosis
		(dry and scaly skin)
		and Keratomalacia
	AFN	Atrophy of lacrymal
	V L	glands and reduction
	DY	in tear secretion
D (Calciferol)/	Promotes intestinal	Rickets in children
Antirachitic vitamin	absorption of calcium	(softness and
	and phosphorus.	deformities of bones
	Formation of teeth	and bow legs and
	and	pigeon chest) and
	bones	Osteomalacia in adults
		(weak and fragile
		bones,
		bent, deformed
		pelvis).
E (Tocopherol) /	Antioxidant	Sterility in animals,
Antisterility	It keeps the skin	Ruptured red blood
vitamin	healthy	cells
	by reduces the	
	process of	
	ageing	



K Anti haemorrahagic	1. Required for the	Defect in blood		
vitamin.	synthesis	clotting		
	of prothrombin in the	called Haemorrhagic		
	liver.	manifestations.		

Water Soluble Vitamins

Vitamins	Functions	Symptoms of Deficiency
B1 (Thiamine)	Involved in	Beriberi: affects
	carbohydrate	muscular, nervous and
	metabolism.	cardiovascular system
	Act as a coenzyme	
B2 (Riboflavin)	Acts as coenzyme in	Inflammation, soreness
	oxidation and	and fissures in the
	reduction	corners of the mouth,
	reactions	lips and tongue.
		Loss of appetite.
		Skin and eye disorder
B3 (Pantothenic	Acts as coenzyme A	Gastrointestinal
acid)	and is essential for the	disorders, anaemia,
	metabolism of fats and	Burning feet
	carbohydrates	syndrome, etc.
B4 (choline)	Precursor for	Fatty liver
2.	acetylcholine	
B5 (Niacin /	Derivatives of	Pellagra (4D
Nicotinic acid)	coenzymes	Syndrome)
		characterized by
		dermatitis, diarrhoea
		and dementia (mental
		deterioration) and
		death.
B6 (Pyridoxine)	Haemoglobin	Dermatitis,
	formation,	convulsions, muscular
	brain, heart and liver	twitching and anaemia
	activities	
B7 (Biotin) / Vit.H	Acts as a coenzyme in	Dermatitis
	synthesis of fat,	
	glycogen and	

		CHENNAL
	amino acids	
B9 (Folic acid)	It acts as a co-enzyme	Megaloblastic anaemia
	for	(large, immature,
	synthesis of nucleic	nucleated RBC in
	acid	blood)
	and essential for	
	growth and formation	
	of RBC	
B12 (Cobalamine)	Promotes DNA	Pernicious anaemia
	synthesis. Necessary	(immature nucleated
	for maturation of RBC	RBC without
	and formation of	haemoglobin). Causes
	myelin sheath.	nervous disorder
C (Ascorbic acid)	Acts as an antioxidant.	Scurvy (Sailor's
	Strengthens the	disease) characterized
	immune	by spongy and
	system.	bleeding gums, falling
	Necessary for healthy	of teeth, fragile bones,
	gums and teeth.	delayed wound
		healing etc Infantile
	CEN	scurvy)

Food adulterants cause harmful effects in the form of headaches, palpitations, allergies, cancers and in addition reduces the quality of food. Common adulterants are addition of citric acid to lemon juice, papaya seeds to pepper, melamine to milk, vanillin for natural vanillin, red dyes to chillis, lead

chromate and lead tetraoxide to turmeric powder, etc.,

Caloric value of carbohydrates, proteins and fats

• We obtain 50% energy from carbohydrates 35% from fats and 15% from proteins. We require about 400 to 500 gm of carbohydrates, 60 to 70 gm of fats and 65 to 75 gm of proteins per day. Balanced diet of each individual will vary according to their age, gender, level of physical activity and others conditions such as pregnancy and lactation.



- Carbohydrates are sugar and starch. These are the major source of cellular fuel which provides energy. The caloric value of carbohydrate is 4.1 calories per gram and its physiological fuel value is 4 Kcal per gram.
- Lipids are fats and derivatives of fats, are also the best reserved food stored in our body which is used for production of energy. Fat has a caloric value of 9.45 Kcal and a physiological fuel value of 9 Kcal per gram.

Many research findings have proven that usage of chemical preservatives and artificial enhancers lead to highly harmful effects. It includes heart ailments, hypertension, infertility, gastrointestinal disorders, early puberty in girls, weakening of bones, damage in organs like kidney and liver, chronic obstructive pulmonary diseases, headache, allergies, asthma, skin rashes and even cancer. Remember that nothing will beat and overtake the taste and safety of homemade foods. "East or west home preparation is the best."

• Proteins are source of amino acids required for growth and repair of body cells. They are stored in the body only to a certain extent; large quantities are excreted as nitrogenous waste. The caloric value and physiological fuel value of one gram of protein are 5.65 Kcal and 4 Kcal respectively. According to ICMR (Indian Council of Medical Research and WHO (World Health Organization), the daily requirement of protein for an average Indian is 1gm per 1 kg body weight.

Nutritional and digestive disorders

• Intestinal tract is more prone to bacterial, viral and parasitic worm infections. This infection may cause inflammation of the inner lining of colon called colitis. The most common symptoms of colitis are rectal bleeding, abdominal cramps, and diarrhoea.

Protein energy malnutrition: (PEM)

• Growing children require more amount of protein for their growth and development. Protein deficient diet during early stage of children



may lead to protein energy malnutrition such as Marasmus and Kwashiorkor. Symptoms are dry skin, pot-belly, oedema in the legs and face, stunted growth, changes in hair colour, weakness and irritability. Marasmus is an acute form of protein malnutrition. This condition is due to a diet with inadequate carbohydrate and protein. Such children are suffer from diarrhoea, body becomes lean and weak (emaciated) with reduced fat and muscle tissue with thin and folded skin.

- Indigestion: It is a digestive disorder in which the food is not properly digested leading to a feeling of fullness of stomach. It may be due to inadequate enzyme secretion, anxiety, food poisoning, over eating, and spicy food.
- Constipation: In this condition, the faeces are retained within the rectum because of irregular bowel movement due to poor intake of fibre in the diet and lack of physical activities.
- Vomiting: It is reverse peristalsis. Harmful substances and contaminated food from stomach are ejected through the mouth. This action is controlled by the vomit centre located in the medulla oblongata. A feeling of nausea precedes vomiting.
- Jaundice: It is the condition in which liver is affected and the defective liver fails to break down haemoglobin and to remove bile pigments from the blood. Deposition of these pigments changes the colour of eye and skin yellow. Sometimes, jaundice is caused due to hepatitis viral infections.
- Liver cirrhosis: Chronic disease of liver results in degeneration and destruction of liver cells resulting in abnormal blood vessel and bile duct leading to the formation of fibrosis. It is also called deserted liver or scarred liver. It is caused due to infection, consumption of poison, malnutrition and alcoholism.
- Gall Stones: Any alteration in the composition of the bile can cause the formation of stones in the gall bladder. The stones are mostly formed of crystallized cholesterol in the bile. The gall stone causes obstruction in the cystic duct, hepatic duct and also hepato-pancreatic duct causing pain, jaundice and pancreatitis.



Appendicitis: It is the inflammation of the vermiform appendix, leading to severe abdominal pain. The treatment involves the removal of appendix by surgery. If treatment is delayed the appendix may rupture and results in infection of the abdomen, called peritonitis.

Hiatus hernia (Diaphragmatic hernia): It is a structural abnormality in which superior part of the stomach protrudes slightly above the diaphragm. The exact cause of hiatus hernias is not known. In some people, injury or other damage may weaken muscle tissue, by applying too much pressure (repeatedly) on the muscles around the stomach while coughing, vomiting, and straining during bowel movement and lifting heavy object. Heart burn is also common in those with a hiatus hernia. In this condition, stomach contents travel back into the oesophagus or even into oral cavity and causes pain in the centre of the chest due to the eroding nature of acidity.

Diarrhoea: It is the most common gastrointestinal disorder worldwide. It is sometimes caused by bacteria or viral infections through food or water. When the colon is infected, the lining of the intestine is damaged by the pathogens, thereby the colon is unable to absorb fluid. The abnormal frequency of bowel movement and increased liquidity of the faecal discharge is known as diarrhoea. Unless the condition is treated, dehydration can occur. Treatment is known as oral hydration therapy. This involves drinking plenty of fluids – sipping small amounts of water at a time to rehydrate the body.


Peptic ulcer: It refers to an eroded area of the tissue lining (mucosa) in the stomach or duodenum. Duodenal ulcer occurs in people in the age group of 25 - 45 years. Gastric ulcer is more common in persons above the age of 50 years. Ulcer is mostly due to infections caused by the bacterium Helicobacter pylori. It may also be caused due to uncontrolled usage of aspirin or certain anti inflammatory drugs. Ulcer may also be caused due to smoking, alcohol, caffeine and psychological stress.

Obesity: It is caused due to the storage of excess of body fat in adipose tissue. It may induce hypertension, atherosclerotic heart disease and diabetes. Obesity may be genetic or due to excess intake of food,



endocrine and metabolic disorders. Degree of obesity is assessed by body mass index (BMI). A normal BMI range for adult is 19- 25; above 25 is considered as obese. BMI is calculated as body weight in Kg, divided by the square of body height in meters. For example, a 50 Kg person with a height of 160 cms would have a BMI of 19.5. That is BMI = 50/1.62 = 19.5

Nobel Prize for the year 2005 was awarded to Robin Warren and Barry Marshall for the discovery of Helicobacter pylori which causes peptic ulcer





12th Zoology

Unit 7 – Human Health and Diseases

Common diseases in human beings

- Disease can be defined as a disorder or malfunction of the mind or body. It involves morphological, physiological and psychological disturbances which may be due to environmental factors or pathogens or genetic anomalies or life style changes. Diseases can be broadly grouped into infectious and non infectious types. Diseases which are transmitted from one person to another are called infectious diseases or communicable diseases. Such disease causing organisms are called pathogens and are transmitted through air, water, food, physical contact and vectors. The disease causing pathogen may be virus, bacteria, fungi, protozoan parasites, helminthic parasites, etc., Infectious diseases are common and everyone suffers from such diseases at some time or the other. Most of the bacterial diseases are curable but all viral diseases are not. Some infectious disease like AIDS may be fatal.
- Non-infectious diseases are not transmitted from an infected person to a healthy person. In origin they may be genetic (cystic fibrosis), nutritional (vitamin deficiency diseases) and degenerative (arthritis, heart attack, stroke). Among non - infectious diseases, cancer is one of the major causes of death.

Bacterial and viral diseases Bacterial diseases

- Though the number of bacterial species is very high, only a few bacteria are associated with human diseases and are called pathogenic bacteria. Such pathogens may emit toxins and affected the body.
- Bacteria spread through air, water or by inhaling the droplets/aerosols or even by sharing utensils, dresses with an infected person. Typhoid fever can be confirmed by Widal test



Viral diseases

- Viruses are the smallest intracellular obligate parasites, which multiply within living cells. Outside the living cells they cannot carry out the characteristics of a living organism. Viruses invade living cells, forcing the cells to create new viruses. The new viruses break out of the cell, killing it and invade other cells in the body, causing diseases in human beings. Rhino viruses cause one of the most infectious human ailment called the "Common cold".
- Viral diseases are generally grouped into four types on the basis of the symptoms produced in the body organs.
- Pneumotropic diseases (respiratory tract infected by influenza)
- Dermotropic diseases (skin and subcutaneous tissues affected by chicken pox and measles)
- Viscerotropic diseases (blood and visceral organs affected by yellow fever and dengue fever)
- Neurotropic diseases (central nervous system affected by rabies and polio).

Bacterial resistance

If an antibiotic is used too often to fight a specific bacterial infection, the bacteria may become resistant to the specific antibiotic. Hence the specific antibiotic can no longer be used to treat the bacterial infection. Some bacteria have developed resistance to many antibiotics. Therefore, infections caused by these bacteria are difficult to be cured. **Risk of bacterial resistance can be reduced by observing the following steps**

- Avoid using antibiotics to treat minor infections that can be taken care by our immune system.
- Do not use an antibiotic to treat viral infections such as common cold or flu.
- Always follow the prescription. Skipping doses or failing to complete the prescription may allow antibiotic resistance to develop.

Bacterial diseases in human beings



S.N o	Diseases	Causative agent	Site of infectio	Mode of transmission	Symptoms
1	Shigellosis (Bacillary dysentery)	Shigella sp.	Intestin e	Food and Water contaminate d by faeces/ faecal oral route	Abdominal pain, dehydratio n, blood and mucus in the stools
2	Bubonic plague (Black death)	Yersinia pestis	Lymph nodes	Rat flea vector – Xenopsylla cheopis	Fever, headache, and swollen lymph nodes.
3	Diphtheria	Corynebacterim diphtheriae	Larynx, skin, nasal and genital passage	Droplet infection	Fever, Sore throat, hoarseness and difficulty in breathing
4	Cholera	Vibrio cholerae	Intestin e	Contaminate d food and water/faecal oral route	Severe diarrhoea and dehydratio n.
5	Tetanus (Lock Jaw)	Clostridium tetani	Spasm of muscles	Through wound infection	Rigidity of jaw muscle, increased heart beat rate and spasm of the muscles of the jaw and face.
6	Typhoid(Enteri c fever)	Salmonella typhi	Intestin e	Through contaminate d	Headache, abdominal discomfort,

				A			
				food	and	fever	and
				water		diarrh	noea.
7	Pneumonia	Streptococcus	Lungs	Droplet		Fever	,
		pneumoniae		infection		cough	ı <i>,</i>
						Painfi	ıl
						breath	ning
						and	brown
						sputu	m
8	Tuberculosis	Mycobacterium	Lungs	Droplet		Thick	
		tuberculosis		infection		muco	purul
						ant	nasal
						discha	arge.

Viral diseases in human beings

S.N	Diseases	Causativ	Site of	Mode of	Symptoms
0		e Agent	infection	transmissio	
			2. //	n	
1	Common	Rhino	Respirator	Droplet	Nasal
	Cold	Viruses	y tract	infection	congestion
			LL		and
		10.			discharge,
	51	UP			sore throat,
	SI				cough and
					headache.
2	Mumps	Mumps	Salivary	Saliva and	Enlargemen
		virus	glands	droplet	t of the
		(RNA		infection	parotid
		virus)			glands.
		Paramyx			
		o virus			
3	Measles	Rubella	Skin and	Droplet	Sore throat,
		virus	respiratory	infection	running
		(RNA			nose, cough
		virus),			and fever.
		Paramyx			Reddish
		o virus			rashes on
					the skin,
					neck and

					CHENNAI
					ears.
4	Viral hepatitis	Hepatitis - B Virus	Liver	Parenterak route, blood transfusion	Liver damage, jaundice, nausea, Yellowish eyes, fever and pain in the abdomen
5	Chicken pox	Varicella Zoster virus (DNA Virus)	Respirator y tract, skin and nervous system	Droplet infection and direct contact	Mild fever with itchy skin, rash and blisters
6	Poliomyelitis	Polio virus (RNA virus)	Intestine, brain, spinal cord	Droplet infection through faecal oral route	Fever, muscular stiffness and weakness, paralysis and respiratory failure
7	Dengue fever (Break bone fever)	Dengue virus or flavi virus (DENV 1-4 virus)	Skin and blood	Mosquito vector Aedes aegypti	Severe flu like illness with a sudden onset of fever and painful headache, muscle and joint pain.
8	Chikunguny a	Alpha virus (Toga virus)	Nervous system	Mosquito vector Aedes aegypti	Fever and joint pain, headache and joint swelling



Nipah virus is a zoonotic virus (transmitted from animals to humans) and also transmitted through contaminated food. In infected people, it causes a range of illness from asymptomatic infection to acute respiratory illness and fatal encephalitis.

Swine flu was first recognised in the 1919 pandemic and still circulates as a seasonal flu virus. Swine flu is caused by the H1N1 virus strain. Symptoms include fever, cough, sore throat, chills, weakness and body aches. Children, pregnant women and the elderly are at risk from severe infection

Protozoan diseases

- About 15 genera of protozoans live as parasites within the human body and cause diseases. Amoebiasis also called amoebic dysentery or amoebic colitis is caused by Entamoeba histolytica, which lives in the human large intestine and feeds on food particles and bacteria. Infective stage of this parasite is the trophozoite, which penetrates the walls of the host intestine (colon) and secretes histolytic enzymes causing ulceration, bleeding, abdominal pain and stools with excess mucus. Symptoms of amoebiasis can range from diarrhoea to dysentery with blood and mucus in the stool. House flies (Musca domestica) acts as a carrier for transmitting the parasite from contaminated faeces and water.
- African sleeping sickness is caused by Trypanosoma species. Trypanosoma is generally transmitted by the blood sucking Tsetse flies. Three species of Trypanosoma cause sleeping sickness in man.
- T. gambiense is transmitted by Glossina palpalis (Tsetse fly) and causes Gambian or Central African sleeping sickness.
- T. rhodesiense is transmitted by Glossina morsitans causing Rhodesian or East African sleeping sickness.



- T. cruzi is transmitted by a bug called Triatoma megista and causes Chagas disease or American trypanosomiasis.
- Kala azar or visceral leishmaniasis is caused by Leishmania donovani, which is transmitted by the vector Phlebotomus (sand fly). Infection may occur in the endothelial cells, bone marrow, liver, lymph glands and blood vessels of the spleen. Symptoms of Kala azar are weight loss, anaemia, fever, enlargement of spleen and liver.
- Malaria is caused by different types of Plasmodium species such as P. vivax, P. ovale, P. malariae and P. falciparum . Plasmodium lives in the RBC of human in its mature condition it is called as trophozoite. It is transmited from one person to another by the bite of the infected female Anopheles mosquito.

Life cycle of Plasmodium

- Plasmodium vivax is a digenic parasite, involving two hosts, man as the secondary host and female Anopheles mosquito as the primary host. The life cycle of Plasmodium involves three phases namely schizogony, gamogony and sporogony
- The parasite first enters the human blood stream through the bite of an infected female Anopheles mosquito. As it feeds, the mosquito injects the saliva containing the sporozoites. The sporozoite within the blood stream immediately enters the hepatic cells of the liver. Further in the liver they undergo multiple asexual fission (schizogony) and produce merozoites. After being released from liver cells, the merozoites penetrate the RBC's.
- Inside the RBC, the merozoite begins to develop as unicellular trophozoites. The trophozoite grows in size and a central vacuole develops pushing them to one side of cytoplasm and becomes the signet ring stage. The trophozoite nucleus then divides asexually to produce the schizont. The large schizont shows yellowish brown pigmented granules called Schuffners granules. The schizont divides and produces mononucleated merozoites. Eventually the erythrocyte lyses, releasing the merozoites and haemozoin toxin into the blood stream to infect other erythrocytes. Lysis of red blood cells results in



cycles of fever and other symptoms. This erythrocytic stage is cyclic and repeats itself approximately every 48 to 72 hours or longer depending on the species of Plasmodium involved. The sudden release of merozoites triggers an attack on the RBCs. Occasionally, merozoites differentiate into macrogametocytes and microgametocytes. When these are ingested by a mosquito, they develop into male and female gametes respectively.

- In the mosquito's gut, the infected erythrocytes lyse and male and female gametes fertilize to form a diploid zygote called ookinete. The ookinete migrates to the mosquito's gut wall and develop into an oocyte. The oocyte undergoes meiosis by a process called sporogony to form sporozoites. These sporozoites migrate to the salivary glands of the mosquito. The cycle is now completed and when the mosquito bites another human host, the sporozoites are injected and the cycle begins a new.
- The pathological changes caused by malaria, affects not only the erythrocytes but also the spleen and other visceral organs. Incubation period of malaria is about 12 days. The early symptoms of malaria are headache, nausea and muscular pain. The classic symptoms first develop with the synchronized release of merozoites, haemozoin toxin and erythrocyte debris into the blood stream resulting in malarial paroxysms shivering chills, high fever followed by sweating. Fever and chills are caused partly by malarial toxins that induce macrophages to release tumour necrosis factor (TNF- α) and interleukin.

S.No		Causative agent	Duration of
	Malaria		Erythrocytic
			Cycle.
1	Tertian, benign	P. Vivax	48 hours
	tertian or vivax		
	malaria		
2	Quartan malaria	P. malariae	72 hours
3	Mild tertian	P. ovale	48 hours
	malaria		
4	Malignant	P. falciparum	36-48 hours

Types of malaria



	tertian	or	
	quotidian		
	malaria		

- Ponds, drainage ditches and other permanent bodies of water can be stocked with fishes such as Gambusia which feed on mosquito larvae. Preparations containing Bacillus thuringiensis can be sprayed to kill the mosquito larvae since it is not toxic to other forms of life. The best protection against malaria is to avoid being bitten by mosquito. People are advised to use mosquito nets, wire gauging of windows and doors to prevent mosquito bites.
- In the 1950's the World Health Organisation (WHO) introduced the Malaria eradication programme. This programme was not successful due to the resistance of Plasmodium to the drugs used to treat it and resistance of mosquito's to DDT and other insecticides.

Malaria vaccine is used to prevent malaria. The only approved vaccine as of 2015 is RTS,S(Mosquirix). It requires four injections and has relatively low efficacy (26–50%). Due to this low efficacy, WHO does not recommend the use of RTS,S vaccine in babies between 6 and 12 weeks of age.

Fungal diseases

- Fungi was recognized as a causative agent of human diseases much earlier than bacteria. Dermatomycosis is a cutaneous infection caused by fungi belonging to the genera Trichophyton, Microsporum and Epidermophyton.
- Ringworm is one of the most common fungal disease in humans. Appearance of dry, scaly lesions on the skin, nails and scalp are the main symptoms of the disease. Heat and moisture help these fungi to grow and makes them to thrive in skin folds such as those in the groin or between the toes. Ringworms of the feet is known as Athlete's foot caused by Tinea pedis. Ringworms are generally acquired from soil or by using clothes, towels and comb used by infected persons.



Helminthic diseases

- Helminthes are mostly endoparasitic in the gut and blood of human beings and cause diseases called helminthiasis. The two most prevalent helminthic diseases are Ascariasis and Filariasis.
- Ascaris is a monogenic parasite and exhibits sexual dimorphism. Ascariasis is a disease caused by the intestinal endoparasite Ascaris lumbricoides commonly called the round worms. It is transmitted through ingestion of embryonated eggs through contaminated food and water. Children playing in contaminated soils are also prone to have a chance of transfer of eggs from hand to mouth. The symptoms of the disease are abdominal pain, vomiting, headache, anaemia, irritability and diarrhoea. A heavy infection can cause nutritional deficiency and severe abdominal pain and causes stunted growth in children. It may also cause enteritis, hepatitis and bronchitis.
- Filariasis is caused by Wuchereria bancrofti, commonly called filarial worm. It is found in the lymph vessels and lymph nodes of man. Wuchereria bancrofti is sexually dimorphic, viviparous and digenic. The life cycle is completed in two hosts, man and the female Culex mosquito The female filarial worm gives rise to juveniles called microfilariae larvae. In the lymph glands, the juveniles develop into adults. The accumulation of the worms block the lymphatic system resulting in inflammation of the lymph nodes.
- In some cases, the obstruction of lymph vessels causes elephantiasis or fi lariasis of the limbs, scrotum and mammary glands

Maintenance of personal and public hygiene

• Hygiene is a set of practices performed to conserve good health. According to the World Health Organization (WHO), hygiene refers to "conditions and practices that help to maintain health and prevent the spread of diseases." Personal hygiene refers to maintaining one's body clean by bathing, washing hands, trimming fingernails, wearing clean clothes and also includes attention to keeping surfaces in the home and workplace, including toilets, bathroom facilities, clean and pathogen-free.



- Our public places teem with infection, contamination and germs. It seems that every surface we touch and the air we breathe are with pollutants and microbes. It's not just the public places that are unclean, but we might be amazed at the number of people who do not wash their hands before taking food, after visiting the restroom, or who sneeze without covering their faces. Many infectious diseases such as typhoid, amoebiasis and ascariasis are transmitted through contaminated food and water.
- Advancement in science and technology provide effective controlling measures for many infectious and non-infectious diseases. The use of vaccines and adopted immunization programmes have helped to eradicate small pox in India. Moreover a large number of infectious diseases like polio, diphtheria, pneumonia and tetanus have been controlled by the use of vaccines and by creating awareness among the people.

Adolescence – drug and alcohol abuse

- Adolescence begins with a period of rapid physical and sexual development called puberty to maturity at 12 to 19 years of age. Adolescence is also a highly dynamic period of psychological and social changes in individuals. Adolescents are vulnerable to group (peer) pressure and many youngsters are pushed into experimenting with drugs and alcohol. Proper education and guidance would enable youth to say no to drugs and alcohol and to follow a healthy life style.
- Alcohol is a psychoactive drug, which acts on the brain, affecting a person's mind and behaviour. It is a depressant, which slows down the activity of the nervous system. The intake of certain drugs for a purpose other than their normal clinical use in an amount and frequency that impair one's physical, physiological and psychological functions is called drug abuse.
- The drugs which are commonly abused include opioids, cannabinoids, coca-alkaloids, barbiturates, amphetamines and LSD.



- Opioids are drugs which bind to specific opioid receptors present in the central nervous system and intestinal tract. Heroin (smack) is chemically diacetyl morphine, which is white, odourless and bitter crystalline compound. It is obtained by acetylation of morphine, which is extracted from flowers of the poppy plant. Morphine is one of the strongest pain killer and is used during surgery. It is the most widely abused narcotic drug which acts as a depressant and slows down body functions.
- Cannobinoids are a group of chemicals obtained from Cannabis sativa, the Indian hemp plant. Natural cannabinoids are the main source of marijuana, ganja, hashish and charas. It interferes in the transport of the neurotransmitter, dopamine and has a potent stimulating action on the CNS, producing increased energy and a sense of euphoria.
- Cocaine is a white powder that is obtained from the leaves of the coca plant, Erythroxylum coca. It is commonly called coke or crack. Cocaine causes serious physical and psychological problems including hallucinations and paranoia. The other plants with hallucinogenic properties are Atropa belladonna and Datura.

Group	Drugs	Effects
Stimulants	Amphetamines,	Accelerates the
	cocaine, nicotine and	activity of the brain
	tobacco	
Depressants	Alcohol, Barbiturates,	Slows down the
	Tranquilizers	activity of the brain
Narcotic/ Analgesics	Opium, Morphine	Act as depressants on
		the Central Nervous
		System
Cannabis	Bhang (Marijuana),	Affects the
	Ganja, Charas	cardiovascular system
Hallucinogens	Lysergic acid	Distorts the way one
	diethylamide (LSD),	sees, hears and feels.
	Phencyclidine	

Classification of drugs



- Drugs like methamphetamine, amphetamines, barbiturates, tranquilizers, Lysergic acid diethylamide (LSD) are normally used as medicine to treat patients with mental illness like depression and insomnia and are often abused.
- Tobacco is smoked, chewed and used as snuff. It increases the carbon monoxide content of blood and reduces the concentration of haem bound oxygen, thus causing oxygen deficiency in the body. Tobacco contains nicotine, carbon monoxide and tars, which cause problems in the heart, lung and nervous system. Adrenal glands are stimulated by nicotine to release adrenaline and nor adrenaline which increases blood pressure and heart beat.

Addiction and dependence

- Addiction is a physical or psychological need to do or take or use certain substance such as alcohol, to the point where it could be harmful to the individual. This addictive behaviour can be personally destructive to a person. Overtime addicts start to lose not only their jobs, homes and money, but also friendship, family relationships and contact with the normal world. Addiction to drugs and alcohol can lead to a psychological attachment to certain effects such as euphoria and temporary feeling of well being.
- Repeated use of drugs and alcohol may affect the tolerance level of the receptors present in the body. These receptors then respond only to highest doses of drugs and alcohol leading to greater intake and addiction. Excessive use of drug and alcohol leads to physical and psychological dependence. When psychological dependence develops, the drug user gets mentally 'hooked on' to the drug. The drug user constantly thinks only about the drug and has continuous uncontrollable craving for it. This state called "euphoria" is characterized by mental and emotional preoccupation with the drug.
- Physical dependence is a state in which the user's body requires a continuous presence of the drug. If the intake of the drug or alcohol is abruptly stopped, he or she would develop withdrawal symptoms. In a sense, the body becomes confused and protests against the absence of the drug. The withdrawal symptoms may range from mild tremors



to convulsions, severe agitation and fits, depressed mood, anxiety, nervousness, restlessness, irritability, insomnia, dryness of throat, etc, depending on the type of drug abuse.

Effects of drugs and alcohol

- Short-term effect appears only for a few minutes after the intake of drugs and alcohol. The abuser feels a false sense of well being and a pleasant drowsiness. Some short term effects are euphoria, pain, dullness of senses, alteration in behaviour, blood pressure, narcosis (deep sleep), nausea and vomiting.
- Drugs and alcohol have long-term effect that lead to serious damages, because of the constant and excessive use. The physical and mental disturbance makes the life of the user unbearable and torturous. For example heavy drinking permanently damages the liver and brain.
- The use of alcohol during adolescence may have long-term effects. Alcohol interferes with the ability of the liver to break down fat. Over time fat accumulation and high levels of alcohol destroy the liver cells and a scar tissue grows in the place of dead cells. This scarring of the liver is called "Liver cirrhosis". Alcohol irritates the stomach lining due to the production of excess acid leading to ulcers. Excessive alcohol use weakens the heart muscle, causing scar tissue to build up in the cardiac muscle fibers. As a result, heavy drinkers have an increased risk of high blood pressure, stroke, coronary artery disease and heart attack. Korsakoff syndrome, a chronic memory disorder is most commonly caused by alcohol misuse.

Alcoholism is the inability to control drinking due to physical and emotional dependence on alcohol. Treatment involves counseling by a healthcare professional. Detoxification programme in a hospital or medical facility is an option for those who need additional assistance. Medications are available to reduce the desire to drink and smoke.



Prevention and control

• It is practically possible to prevent some one from using drugs and alcohol. Here are some ways that help to prevent drug and alcohol abuse.

Effectively dealing with peer pressure

The biggest reason for teens to start on drugs is due to their friends / peer groups imposing pressure on them. Hence, it is important to have a better group of friends to avoid such harmful drugs and alcohol.

Seeking help from parents and peers

Help from parents and peer group should be sought immediately so that they can be guided appropriately. Help may even be sought from close and trusted friends. Getting proper advice to sort out their problems would help the young to vent their feelings of anxiety and guilt.

Education and counselling

Education and counselling create positive attitude to deal with many problems and to accept disappointments in life.

Looking for danger signs

Teachers and parents need to look for sign that indicate tendency to go in for addiction.

Seeking professional and medical assistance

Assistance is available in the form of highly qualified psychologists, psychiatrists and de-addiction and rehabilitation programmes to help individuals to overcome their problems.

Mental health – Depression

 Mental health is a state of well being of the mind, with self esteem. Self esteem means liking yourself and being able to stand up for what you believe is right. Positive mental health is an important part of wellness. A mentally healthy person reflects a good personality. Activities of mentally healthy people are always appreciated and



rewarded by the society as these persons are creative as well as cooperative with others. Mental health improves the quality of life.

• Depression is a common mental disorder that causes people to experience depressed mood, loss of interest or pleasure, feelings of guilt or low self-worth, disturbed sleep poor appetite, low energy and poor concentration.

Alcoholic Anonymous

Alcoholic anonymous was started in 1935 by a businessman and a doctor who had been a "hopeless drunk" for many years. After the men helped each other to stop drinking and to stay sober, they then founded the alcoholic anonymous to help other alcoholics. Since that time alcoholic anonymous has spread throughout the world.

Signs and symptoms of mental depression

- Loss of self confidence and self esteem
- > Anxiety
- Not being able to enjoy things that are usually pleasurable or interesting.
- Lifestyle changes like exercise, meditation, yoga and healthy food habits can help to be relieved from depression. Exercise stimulates the body to produce serotonin and endorphins, which are neurotransmitters that suppress depression. Practicing exercise in daily life creates a positive attitude .

Participating in an exercise programme can:

- Increase self-esteem
- Boost self-confidence
- Create a sense of empowerment
- Enhance social connections and relationships
- Brain is one of the most metabolically active part of the body and needs a steady stream of nutrients to function. A poor diet may not provide the nutrients for a healthy body and may provoke symptoms of anxiety and depression.



Lifestyle disorders in human beings

- The old saying that "health is wealth" is truly applicable to human beings. With the changes in life style, there are many emerging medical conditions and diseases that are reducing human longevity. Life style disorder result due to activities involving smoking, alcohol and drug abuse, consuming high fat diet, lack of exercise or living a latent life.
- The World Health Organization (WHO) in its report cautions a slow moving public health disaster due to life style disorders in the form of non-communicable diseases like diabetes, cardiovascular and lung diseases. WHO believes that not thousands but millions of people die every year within the age group of thirty to sixty due to life style related disorders.
- The following facts will help in better understanding of life style disorders.
- Life style disorder causes cardiovascular diseases resulting in 31 percent of global deaths.
- The sedentary life style also causes deficiency of vitamins such as vitamin D resulting in fatigue, tiredness, back pain, depression, bone loss, muscle pain, etc,
- Life style disorder also includes social isolation resulting in age related problems.
- Eating junk foods that have high caloric values, rich in carbohydrates and fat can lead to obesity and early health issues.
- Consumption of processed and packaged food, which lacks in fiber may result in constipation.
- Several people today complain of irritable bowel syndrome with stomach discomfort or pain and trouble with bowel movements, causing diarrhoea. The main cause of irritable bowel syndrome is stress and other illnesses.




Life style modifications

- Avoid eating junk food and foods that have preservatives and colouring agents.
- Physical exercises such as brisk walking and yoga can be done regularly.
- Following medical advice, if any health problems in addition to life style disorders.

- > To avoid smoking drugs and drinking alcohol.
- > To follow a healthy balanced diet rich in vitamins and proteins.
- > 7 8 hours of sleep every day is required.

TUD





Chapter 8 – Immunology

8. Immunology

• In the previous chapter, we have studied in detail the various infections which cause diseases in human beings. In this chapter, we shall discuss how our body protects us from these infections by the effective mechanism of the immune system.

Basic concepts of immunology

- Immunology is the study of immune system. This system protects an individual from various infective agents. It refers to all the mechanisms used by the body for protection from environmental agents that are foreign to the body.
- When the immune system does not function efficiently in an individual, it leads to infection causing disease. The overall ability of body to fight against the disease causing pathogen is called immunity. It is also called disease resistance and the lack of immunity is known as susceptibility. Immunity is highly specific.
- Normally many of the responses of the immune system initiate the destruction and elimination of invading organisms and any toxic molecules produced by them. These immune reactions are destructive in nature and are made in response only to molecules that are foreign to the host and not to those of host itself. This ability to distinguish foreign molecules from self is another fundamental feature of the immune system. However, occasionally, it fails to make its distinction and reacts destructively against the host's own molecules; such autoimmune diseases can be fatal to the organism.
- Almost all the macromolecules e.g. proteins, polysaccharides, nucleic acids, etc., as long as they are foreign to recipient organism can induce immune response. Any substance capable of eliciting immune response is called an ANTIGEN (Antibody Generator). There are two



broad classes of immunity responses namely, innate immunity and acquired immunity (Fig. 8.1).




Fig. 8.1 Immune system

Type of innate Immunity	Mechanism	
1. Anatomical barriers		
• Skin	• Prevents the entry of microbes. Its acidic environment (pH 3-5) retards the growth of microbes.	
Mucus Membrane	• Mucus entraps foreign microorganisms and competes with microbes for attachment.	
2. Physiological barriers		
• Temperature	 Normal body temperature inhibits the growth of pathogens. Fever also inhibits the growth of pathogens. 	
• Low pH	 Acidity of gastric secretions (HCl) kills most ingested microbes. 	
• Chemical mediators	 Lysozyme acts as antibacterial agent and cleaves the bacterial cell wall. Interferon's induce antiviral state in the uninfected cells. Complementary substances produced from leucocytes lyse the pathogenic microbes or facilitate phagocytosis. 	
3. Phagocytic bazrriers	Specialized cells (Monocytes, neutrophils,	

76 | P a g e APPOLO STUDY CENTRE PH: 044-24339436, 42867555, 9840226187

	CHENNAL		
	tissue macrophages) phagocytose, and		
	digest whole micro -organisms.		
4. Inflammatory barriers	Tissue damage and infection induce leakage		
	of vascular fluid, containing chemotactic		
	signals like serotonin, histamine and		
	prostaglandins. They influx the phagocytic		
	cells into the affected area. This		
	phenomenon is called diapedesis.		

8.2 Innate immunity

• Innate immunity is the natural phenomenon of resistance to infection which an individual possesses right from the birth. The innate defense mechanisms are non-specific in the sense that they are effective against a wide range of potentially infectious agents. It is otherwise known as non-specific immunity or natural immunity. A number of innate defense mechanisms are operative non-specifically against a large number of microorganisms as shown in the Table 8.1 ENTR and Fig. 8.2.

Acquired immunity

- The immunity that an individual acquires after birth is known as acquired immunity. It is the body's resistance to a specific pathogen.
- The unique features of acquired immunity are antigenic specificity, diversity, recognition of self and non-self and immunological memory.

Components of acquired immunity

Acquired immunity has two components – cell mediated immunity (CMI) and antibody mediated immunity or humeral immunity.

1. Cell mediated immunity

• When pathogens are destroyed by cells without producing antibodies, then it is known as cell mediated immune response or cell mediated immunity. This is brought about by T cells, macrophages and natural killer cells.



2. Antibody mediated immunity or humoral immunity

• When pathogens are destroyed by the production of antibodies, then it is known as antibody mediated or humoral immunity. This is brought about by B cells with the help of antigen presenting cells and T helper cells. Antibody production is the characteristic feature of vertebrates only.

Types of acquired immunity

• Acquired immunity may be active immunity or passive immunity (Table 8.2). The immunological resistance developed by the organisms through the production of antibodies in their body is called active immunity. Active immunity is acquired through the use of a person's immune responses, which lead to the development of memory cells. Active immunity results from an infection or an immunization. Passive immunity does not require the body to produce antibodies to antigens. The antibodies are introduced from outside into the organism. Thus, passive immunity is acquired without the activation of a person's immune response, and therefore there is no memory.

The process of production of blood cells in the bone marrow is called haematopoiesis.

S. No	Active Immunity	Passive immunity	
1.	Active immunity is produced	Passive Immunity is received	
	actively by host's immune	passively and there is no active	
	system.	host participation.	
2.	It is produced due to contact	It is produced due to antibodies	
	with pathogen or bu its	obtained from outside.	
	antigen.		
3.	It is durable and effective in	It is transient and les effective.	
	protection		
4.	Immunological memory is	No memory.	
	present.		
5.	Booster effect on subsequent	-	
	does is possible.	effective.	

78 | P a g e APPOLO STUDY CENTRE PH: 044-24339436, 42867555, 9840226187



6.	Immunity is effective	only	Immunity	develops
	after a short period.	-	immediately	7

Immune responses

• The immune responses may be primary or secondary (Table 8.3).

Primary immune response

• The primary immune response occurs when a pathogen comes in contact with the immune system for the first time. During this, the immune system has to learn to recognize the antigen, produce antibody against it and eventually produce memory lymphocytes. The primary immune response is slow and short – lived.

S.	Primary Immune Response	Secondary Immune Response	
No			
1.	It occurs as a result of primary	It occurs as a result of second	
	contact with an antigen.	and subsequent contacts with	
		the same antigen.	
2.	Antibody level reaches peak	Antibody level reaches peak in	
	in 7 to 10 days.	3 to 5 days.	
3.	Prolonged periods is required	It establishes immunity in a	
	TUP	short time.	
4.	There is rapid decline in	Antibody level remains high	
	antibody level.	for longer period.	
5.	It appears mainly in the	It appears mainly in the bone	
	lymph nodes and spleen.	marrow, followed by the spleen	
		and lymph nodes.	

Secondary immune response

• The secondary immune response occurs when a person is exposed to the same antigen again. During this, time immunological memory has been established and the immune system can start producing antibodies immediately. Within hours after recognition of the antigen, a new army of plasma cells are generated. Within 2 to 3 days, the antibody concentration in the blood rises steeply to reach much



higher level than primary response. This is also called as "booster response".

Lymphoid organs

• Immune system of an organism consists of several structurally and functionally different organs and tissues that are widely dispersed in the body. The organs involved in the organ, maturation and proliferation of lymphocytes are called lymphoid organs (Fig.8.3). Based on their functions, they are classified into primary or central lymphoid organs and secondary or peripheral lymphoid organs trap antigens and make it available for mature lymphocytes, which can effectively fight against these antigens.

Primary lymphoid organs

• Bursa of Fabricius of birds, bone marrow and thymus gland of mammals constitute the primary lymphoid organs involved in the production and early selection of lymphocytes. These lymphocytes become dedicated to a particular antigenic specificity. Only when the lymphocytes mature in the primary lymphoidal organs, they become immunocompetent cells. In mammals, B cell maturation occurs in the bone marrow and T cells maturation occurs in the thymus.

Thymus

The thymus is a flat and bilobed organ located behind the sternun, above the heart. Each lobe of the thymus contains numerous lobules, separated from each other by connective tissue called septa. Each compartments, lobule is differentiated into outer two the compartment or outer cortex, is densely packed with immature T cells called thymocytes, whereas the inner compartment or medulla is sparsely populated with thymocytes. One of its main secretions is the hormone thymosin. It stimulates the T cell to become mature and immune competent. By the early teens, the thymus begins to atrophy and is replaced by adipose tissue (Fig. 8.4). Thus thymus is most active during the neonatal and pre-adolescent periods.



Bone marrow

• Bone marrow is a lymphoid tissue found within the spongy portion of the bone. Bone marrow contains stem cells known as haematopoietic cells. These cells have the potential to multiply through cell division and either remain as stem cells or differentiate and mature into different kinds of blood cells.

Secondary or peripheral lymphoid organs

• In secondary or peripheral lymphoid organs, antigen is localized so that it can be effectively exposed to mature lymphocytes. The best examples are lymph nodes, appendix, Peyer's patches of gastrointestinal tract, tonsils, adenoids, spleen, MALT (Mucosal-Associated Lymphoid Tissue), GALT (Gut-Associated Lymphoid Tissue), BALT (Bronchial/Tracheal-Associated Lymphoid Tissue).

Peyer's patches are oval-shaped areas of thickened tissue that are embedded in the mucus-secreting lining of the small intestine of humans and other vertebrate animals. Peyer's patches contain a variety of immune cells, including macrophages, dendritic cells, T cells, and B cells.

The tonsils (palatine tonsils) are a pair of soft tissue masses located at the back of the throat (pharynx). The tonsils are part of the lymphatic system, which help to fight infections. They stop invading germs including bacteria and viruses.

Spleen is a secondary lymphoid organ located in the upper part of the abdominal cavity close to the diaphragm. Spleen contains B and T cells. It brings humoral and cell mediated immunity.

• The adenoids are glands located in the roof of the mouth, behind the soft palate where the nose connects to the throat. The adenoids produce antibodies that help to fight infections. Typically, the adenoids shrink during adolescence and may disappear by adulthood.



Lymph node

- Lymph node is a small bean-shaped structure and is part of the body's immune system. It is the first one to encounter the antigen that enters the tissue spaces. Lymph nodes filter and trap substances that travel through the lymphatic fluid. They are packed tightly with white blood cells, namely lymphocytes and macrophages. There are hundreds of lymph nodes found throughout the body. They are connected to one another by lymph vessels. Lymph is a clear, transparent, colourless, mobile and extracellular fluid connective tissue. As the lymph percolates through the lymph node, the particulate antigen brought in by the lymph will be trapped by the phagocytic cells, follicular and inter digitating dendritic cells.
- Lymph node has three zones (Fig. 8.5). They are the cortex, paracortex and medulla. The outer most layer of the lymph node is called cortex, which consists of B-lymphocytes, macrophages, and follicular dendritic cells. The paracortex zone is beneath the cortex, which is richly populated by T lymphocytes and interdigitating dendritic cells. The inner most zone is called the medulla which is sparsely populated by lymphocytes, but many of them are plasma cells, which actively secrete antibody molecules. As the lymph enters, it slowly percolates through the cortex, paracortex and medulla, giving sufficient chance for the phagocytic cells and dendritic cells to trap the antigen brought by the lymph. The lymph leaving a node carries enriched antibodies secreted by the medullary plasma cells against the antigens that enter the lymph node. Sometimes visible swelling of lymph nodes occurs due to active immune response and increased concentration of lymphocytes. Thus swollen lymph nodes may signal an infection. There are several groups of lymph nodes are found in the neck, under the chin, in the armpits and in the groin.

The mucosa – associated lymphoid tissue (MALT) is a diffuse system of small concentrations of lymphoid tissue in the alimentary, respiratory and urino-genital tracts. MALT is populated by lymphocytes such as T and B cells, as well as plasma cells and macrophages, each of which is well situated to encounter antigens passing through the mucosal epithelium.



Gut-associated lymphoid tissue (GALT) is a component of the mucosaassociated lymphoid tissue (MALT) which worked in the immune system to protect the body from invasion in the gut.

Bronchus Associated Lymphoid Tissues **(BALT)** also a component of MALT is made of lymphoid tissue (tonsils, lymph nodes, lymph follicles) is found in the respiratory mucosae from the nasal cavities to the lungs.

Cells of the immune system

• The immune system is composed of many interdependent cells that protect the body from microbial infections and the growth of tumour cells. The cellular composition of adult human blood is given in Table8. 4.

Cell type	Number of cells per μ1	Approximate percentage		
Red blood	4200,000- 6500,000	<u> </u>		
White blood cells				
i Agranulocytes				
 Lymphocytes 	1500 - 4000	20 - 30		
Monocytes	200 - 950	2 – 7		
ii Granulocytes	•			
Neutrophils	2000 - 7000	50-70		
Basophils	50 - 100	<1		
Eosinophils	40 - 500	2 – 5		
Platelets	150,000 - 500,000			

• All these cells are derived from pluripotent haematopoetic stem cells. Each stem cell had the capacity to produce RBC, WBC and platelets. The only cells capable of specifically recognising and producing an immune response are the lymphocytes. The other types of white blood cells play an important role in non-specific immune response, antigen presentation and cytokine production.



Lymphocytes

- About 20-30% of the white blood cells are lymphocytes. They have a large nucleus filling most of the cell, surrounded by a little cytoplasm. The two main types of lymphocytes are B and T lymphocytes. Both these are produced in the bone marrow. B lymphocytes (B cells) stay in the bone marrow until they are mature. Then they circulate around the body. Some remain in the blood, while others accumulate in the lymph nodes and spleen. T lymphocytes leave the bone marrow and mature in the thymus gland. Once mature, T cells also accumulate in the same areas of the body as B cells. Lymphocytes have receptor proteins on their surface. When receptors on a B cell bind with an antigen, the B cell becomes activated and divides rapidly to produce plasma cells. The plasma cells produce antibodies. Some B cells do not produce antibodies but become memory cells. These cells are responsible for secondary immune response. T lymphocytes do not produce antibodies. They recognize antigen-presenting cells and destroy them. The two important types of T cells are Helper T cells and Killer T cells. Helper T cells release a chemical called cytokine which activates B cells. Killer cells move around the body and destroy cells which are damaged or infected (Fig. 8.6).
- Apart from these cells neutrophils and monocytes destroy foreign cells by phagocytosis. Monocytes when they mature into large cells, they are called macrophages which perform phagocytosis on any foreign organism.

Dendritic cells are called so because it's covered with long, thin membrane extensions that resemble dendrites of nerve cells. These cells present the antigen to T-helper cells. Four types of dendritic cells are known. They are Langerhans, interstitial cells, myeloid and lymphoid cells.

Antigens

• The term antigen (Ag) is used in two senses, the first to describe a molecule which generates an immune response and the second, a molecule which reacts with antibodies. In general antigens are large, complex molecular substances that can induce a detectable immune



response. Thus an antigen is a substance that is specific to an antibody or a T-cell receptor and is often used as a synonym for immunogen.

The histocompatibility antigens are cell surface antigens that induce an immune response leading to rejection of allografts.

• An immunogen is a substance capable of initiating an immune response. Haptens are substance that are non-immunogenic but can react with the products of a specific immune response. Substances that can enhance the immune response to an antigen are called adjuvants. Epitope is an antigenic determinant and is the active part of an antigen. A paratope is the antigen – binding site and is a part of an antibody which recognizes and binds to an antigen.

Antigenicity is the property of a substance (antigen) that allows it to react with the products of the specific immune response.

Types of antigens

• On the basis of origin, antigens are classified into exogenous antigens and endogenous antigens. The antigens which enter the host from the outside in the form of microorganisms, pollens, drugs, or pollutants are called exogenous antigens. The antigens which are formed within the individual are endogenous antigens. The best examples are blood group antigens.

Antibodies

• Antibodies are immunoglobulin (Ig) protein molecules synthesized on exposure to antigen that can combine specifically with the antigen. Whenever pathogens enter our body, the B-lymphocytes produce an army of proteins called antibodies to fight with them. Thus, they are secreted in response to an antigen (Ag) by the effect of B cells called plasma cells. The antibodies are classified into five major categories, based on their physiological and biochemical properties. They are IgG (gamma), IgM (mu), IgA (alpha), IgD (delta) and IgE (epsilon).



- In the 1950s, experiments by Porter and Edelman revealed the basic structure of the immunoglobulin. An antibody molecule is Y shaped structure that comprises of four polypeptide chains, two identical light chains (L) of molecular weight 25,000 Da (approximately 214 amino acids) and two identical heavy chains (H) of molecular weight 50,000 Da (approximately 450 amino acids). The polypeptide chains are linked together by di-sulphide (S-S) bonds. One light chain is attached to each heavy chain and two heavy chains are attached to each other to form a Y shaped (Fig. 8.7) structure. Hence, an antibody is represented by H2 L2. The heavy chains have a flexible hinge region at their approximate middles.
- Each chain (L and H) has two terminals. They are C terminal (Carboxyl) and amino or N-terminal. Each chain (L and H) has two regions. They have variable (V) region at one end and a much larger constant (C) region at the other end. Antibodies responding to different antigens have very different (V) regions but their (C) regions are the same in all antibodies. In each arm of the monomer antibody, the (V) regions of the heavy and light chains combines to form an antigen binding site shaped to 'fit' a specific antigenic determinant. Consequently each antibody monomer has two such antigen binding regions. The (C) regions that forms the stem of the antibody monomer determine the antibody class and serve common functions in all antibodies. The functions of immunoglobulin are agglutination, precipitation, opsonisation, neutralization etc.,

Antigen and antibody interaction

• The reaction between an antigen and antibody is the basis for humoral immunity or antibody mediated immunity. The reaction between antigen and antibody occurs in three stages. During the first stage, the reaction involves the formation of antigen - antibody complex. The next stage leads to visible events like precipitation, agglutination, etc,. The final stage includes destruction of antigen or its neutralization (Fig. 8.8).



Binding force of antigen - antibody reaction

- The binding force between antigen and antibody is due to three factors. They are closeness between antigen and antibody, non-covalent bonds or intermolecular forces and affinity of antibody. When antigen and antibody are closely fitted, the strength of binding is great. When they are apart binding strength is low. The bonds that hold the antigen to the antibody combining site are all non-covalent in nature. These include hydrogen bonds, electrostatic bonds, Van der Waals forces and hydrophobic bonds. Antibody affinity is the strength of the reaction between a single antigenic determinant and a single combining site on the antibody.
- The chief application of antigen antibody reactions are to determine blood groups for transfusion, to study serological ascertainment of exposure to infectious agents, to develop immunoassays for the quantification of various substances, to detect the presence or absence of protein in serum and to determine the characteristics of certain immunodeficiency diseases.

Different types of antigen and antibody reactions

- The reaction between soluble antigen and antibody leads to visible precipitate formation, which is called precipitin reaction. Antibodies that bring about precipitate formation on reacting with antigens are called as precipitins.
- Whenever a particulate antigen interacts with its antibody, it would result in clumping or agglutination of the particulate antigen, which is called agglutination reaction. The antibody involved in bringing about agglutination reaction is called agglutinin.
- Opsonisation or enhanced attachment is the process by which a pathogen is marked of ingestion and destruction by a phagocyte. Opsonisation involves the binding of an opsonin i.e., antibody, to a receptor on the pathogen's cell membrane. After opsonin binds to the membrane, phagocytes are attracted to the pathogen. So, opsonisation is a process in which pathogens are coated with a substance called an opsonin, marking the pathogen out for



destruction by the immune system. This results in a much more efficient phagocytosis.

• The neutralization reactions are the reactions of antigen-antibody that involve the elimination of harmful effects of bacterial exotoxins or a virus by specific antibodies. These neutralizing substances i.e., antibodies are known as antitoxins. This specific antibody is produced by a host cell in response to a bacterial exotoxin or corresponding toxoid (inactivated toxin).

Vaccines

- A vaccine is a biological preparation that provides active acquired immunity to a particular disease and resembles a microorganism and is often made from weakened of attenuated or killed forms of the microbes, their toxins, or one of its surface proteins. Vaccines "teach "our body how to defend itself when viruses or bacteria, invade it/ Vaccines deliver only very little amounts of inactivated or weakened viruses or bacteria, or parts of them. This allows the immune system to recognize the organism without actually experiencing the diseases. Some vaccines need to be given more than once (i.e., a 'booster' vaccination) to make sure the immune system can overcome a real infection in the future.
- First generation vaccine is further subdivided into live attenuated vaccine, killed vaccine and toxoids (Fig. 8.9). Live attenuated vaccines use the weakened (attenuated), aged, less virulent form of the virus. E.g. Measles, mumps and rubella (MMR) vaccine and the Varicella (chickenpox) vaccine, Killed (inactivated) vaccines are killed or inactivated by heat and other methods. E.g. Salk's polio vaccine. Toxoid vaccines contain a toxin or chemical secreted by the bacteria or virus. They make us immune to the harmful effects of the infection, instead of to the infection itself. E.g. DPT vaccine (Diphtheria, Pertussis and Tetanus).
- Second generation vaccine contains the pure surface antigen of the pathogen. E.g. Hepatitis-B vaccine. Third generation vaccine contains the purest and the highest potency vaccines which are synthetic in



generation. The latest revolution in vaccine is DNA vaccine or recombinant vaccine (Refer Chapter- 10 for details).

Vaccino therapy is the method of use of vaccine from treatment of disease. Dr. Edward Jenner prepared first vaccine for small pox in 1796. Polio vaccine was developed by Dr. Jonas Salk (vaccine consists of inactivated microorganism) and Dr. Albert Sabin (live attenuated oral polia caccine). Louis Pasteur (1885) discovered vaccine against rabies, anthrax and cholera. BCG vaccine was developed by Calmette and Guerin against tuberculosis in France in the Year 1908.

Vaccination and immunization

 "Vaccination is the process of administrating a vaccine into the body or the act of introducing a vaccine into the body to produce immunity to a specific disease." Immunization is the process of the body building up immunity to a particular disease. Immunization describes the actual changes in the body after receiving a vaccine. Vaccines work by fighting the pathogen and then recording it in their memory system to ensure that the next time this pathogen enters the body, it is eliminated far quickly. Once, the body is able to fight against the disease, it is believed to have built the immunity for it, also known as the body being immunized against the disease.

Hypersentivity - Overactive Immune Response

• Some of the individuals are very sensitive to some particles present in the environment. The exaggerated response of the immune system to certain antigens present in the environment is called allergy (alloaltered, erg-reaction). The substances to which such an immune response is produced are called allergens. An allergen is an antigen that causes an allergic reaction. Allergic reactions begin within few seconds after the contact with the allergen and last about half an hour. The common examples of allergens are mites in dust, pollens and some proteins in insect venom. Hay fever and asthma are some common examples of allergy. Symptoms of allergic reactions include sneezing, watery eyes, running nose and difficulty in breathing. Allergy is a form of over active immune response mediated by IgE and mast cells. It can also be due to the release of chemicals like histamine and serotonin from the mast cells. Anaphylaxis is the



classical immediate hypersensitivity reaction. It is a sudden, systematic, severe and immediate hypersensitivity reaction occurring as a result of rapid generalized mast-cell degranulation.

Immunodeficiency disease – Ineffective immune response – AIDS

• Immunodeficiency results from the failure of one or more components of the immune system. Primary immune deficiencies are caused by genetic developmental defects. Secondary immune deficiencies arise due to various reasons like radiation, use of cytolytic and immunosuppressive drugs and infections. AIDS is an acronym for Acquired Immuno Deficiency Syndrome. It is the deficiency of immune system, acquired during the life time of an individual indicating that it is not a congenital disease. AIDS is caused by Human Immuno Deficiency Virus (HIV). It selectively infects helper T cells. The infected helper T cells will not stimulate antibody production by B-cells resulting in loss of natural defence against viral infection. On the basis of genetic characteristics and differences in the viral antigens, HIV is classified into the types 1 and 2 (HIV-1, HIV-2). GEN

Structure of HIV

The human immunodeficiency virus belongs to the genus Lentivirus. When observed under the electron microscope, HIV is seen as a spherical virus, 100-120 nm in diameter, containing a dense core surrounded by a lipoprotein envelope. The envelope has glycoprotein (gp) spikes termed gp 41 and gp 120. At the core, there are two large single stranded RNA. Attached to the RNA are molecules of reverse transcriptase. It also contains enzymes like protease and ribonuclease. The core is covered by a capsid made of proteins. This is followed by another layer of matrix proteins as shown in the Fig 8.10.

HIV Transmission

• The HIV is often located within the cells especially in macrophages. HIV can survive for 1.5 days inside a cell but only about 6 hours outside a cell. Routes of HIV transmission include unsafe sexual contact, blood-contaminated needles, organ transplants, blood



transfusion and vertical transmission from HIV infected mother to child. HIV is not transmitted by insects or by casual contact.

- After getting into the body of the person, the virus enters into macrophages where RNA genome of the virus replicates to form viral DNA with the help of the enzyme reverse transcriptase. This viral DNA gets incorporated into the DNA of host cells and directs the infected cells to produce viral particles. The macrophages continue to produce virus and in this way acts like a HIV factory. Simultaneously, HIV enters into helper T-lymphocytes, replicates and produces progeny viruses. The progeny viruses released in the blood attack other helper T-lymphocytes. This is repeated, leading to a progressive decrease in the number of helper T lymphocytes in the body of the infected person. During this period, the person suffers from bouts of fever, diarrhoea and weight loss. Due to decrease in the number of helper T lymphocytes, the person starts suffering from infections and becomes immune deficient and unable to protect against any infection.
- A simple blood test is available that can determine whether the person has been infected with HIV. The ELISA test (Enzyme Linked Immune Sorbent Assay) detects the presence of HIV antibodies. It is a preliminary test. Western blot test is more reliable and a confirmatory test. It detects the viral core proteins. If both tests detect the presence of the antibodies, the person is considered to be HIV positive.
- AIDS has no cure. Prevention of AIDS is the best option. Advocating safe sex and promoting regular check-up, safe blood for transfusion, use of disposable needles, use of condoms during sexual contact, prevention of drug abuse, AIDS awareness programme by NACO (National AIDS Control Organisation), NGOs (Non-Governmental Organisations) and WHO are to prevent the spreading of AIDS.

Autoimmune diseases

• Autoimmunity is due to an abnormal immune response in which the immune system fails to properly distinguish between self and non-self and attacks its own body. Our body produces antibodies (auto antibodies) and cytotoxic T cells that destroy our own tissues. If a disease-state results, it is referred to as auto-immune disease. Thus,



autoimmunity is a misdirected immune response. Autoimmunity is evidenced by the presence of auto antibodies and T cells that are reactive with host antigens. When the cells act as antigens in the same body, they are called auto antigens.

• Autoimmune diseases in human can be divided into two broad categories, namely organ-specific and non-organ-specific (systemic) autoimmune diseases. In organ-specific disease, the autoimmune process is directed mostly against one organ. The autoantibodies may block the functions performed by the organs. Examples include Hashimoto's thyroiditis, Graves' disease (thyroid gland) and Addison's disease (adrenal glands). In non-organ specific (systemic) disorders, autoimmune activity is widely spread throughout the body. Rheumatoid arthritis and multiple sclerosis are example for systemic disorder.

Tumour immunology

- A tumour or neoplasm is a group of cells whose growth has gone unchecked. When a tumour continues to grow and invades healthy tissue, it is called cancer. They spread to other parts of the body from the tumour and give rise to secondary tumour. This is known as metastasis. Tumour may be benign or malignant depending on its characteristics. Benign or non-cancerous tissues are capable of indefinite growth and do not invade other body parts. In the malignant tumour, the cells grow indefinitely, detach and migrate into healthy surrounding tissues.
- In normal cells, cell growth and differentiation is highly controlled and regulated. But in cancer cells, there is breakdown of this regulatory mechanism. Normal cells show a property called contact inhibition, which inhibits uncontrolled growth. Cancer cells do not have this property. As a result, cancerous cells divide continuously giving rise to mass of tissues called tumours (Table 8.5).
- When a cell undergoes malignant transformation, it acquires new surface antigen and may also lose some normal antigens. These antigens are present on the membranes of malignant cells and they induce an immune response. Both humoral and cellular responses



can be observed in malignancy. Cancer cells can avoid immune detection as they are not foreign bodies but are abnormally functioning body cells. This makes them difficult to treat.

• The concept of immunological surveillance postulates that the primary function of the immune system is to "seek and destroy" malignant cells that arise by somatic mutation. The efficiency of the surveillance mechanism reduces either as a result of ageing or due to congenital or acquired immune deficiencies, leads to increased incidence of cancer. Thus, if immunological surveillance is effective, cancer should not occur. The development of tumour represents a lapse in surveillance.

Immunotherapy of cancer

• Immunotherapy also called biological therapy uses substances made by the body or in a laboratory (monoclonal antibodies) to improve or to resist the immune system function. Different approaches have been attempted in the immunotherapy of cancer. Immunotherapy appears to be important in getting rid of the residual malignant cells after the gross tumour has been removed. The best results in the treatment of cancer is to follow an integrated approach to therapy, combining surgery, radiotherapy, chemotherapy and immunotherapy.

Scope of Immunology

The younger graduates in this field can find number of employment opportunities in Government as well as private hospitals. The scope of the immunology is immunotherapy, microbial immunology, clinical immunology, cellular immunology, allergy and immunology, translational immunology, transplantation immunology, neuroinflammatory disorders, tumour immunology, vaccine immunology, inflammatory disorders, ocular immunology and inflammation.

Summary

• Immunology deals with a study of the immune system. The immune system recognises and eliminates the invaders, and the ability of the body to overcome the pathogen is called immunity. Immunity is classified into innate immunity and acquired immunity. Acquired



immunity is further classified into cell mediated immunity and antibody mediated immunity as its components. Acquired immunity may be active or passive immunity. Immune response is the body's response to pathogens and it may be primary or secondary. The organs involved in the origin, maturation and proliferation of lymphocytes are called lymphoid organs. Thymus, bone marrow are primary lymphoid organs. The secondary lymphoid organs are lymph node, MALT, GALT and BALT.

• An antigen is a substance that is specific to an antibody. An immunogen is a substance capable of initiating an immune response. Haptens are substance that are non-immunogenic but can react with the product of a specific immune response. Substances that can enhance the immune response to an antigen are called adjuvants. An epitope is also known as antigenic determinant and is the active part of the antigen. A paratope is the part of antibody. Precipitation, agglutination, neutralization, opsonisation etc. are the different types of antigen and antibody reaction. A vaccine is biological preparation that provides active acquired immunity. The malfunctioning of immune system leads to hypersensitivity, immunodeficiency or autoimmune diseases. A tumour or neoplasm is a group of cells whose growth has gone unchecked. The best results in the treatment of cancer is achieved by an integrated approach to therapy, surgery, radiotherapy, chemotherapy and immunotherapy.



12th zoology Unit - 9. Microbes in Human Welfare

Microbes such as bacteria, fungi, protozoa, certain algae, viruses, viroids and prions are some of the major components of the biological system on Earth. Several microorganisms are beneficial and contribute to human welfare. Microbes are presenteverywhere – in soil, water, air and within bodies of animals and plants. Microbes like bacteria and fungi can be grown on nutritive media to form colonies which can be visibly seen. Some of the microbes useful to human welfare are discussed here.

Microbes in household products

- In every-day life, microbes and their products are used in the preparation of idli, dosa, cheese, curd, yogurt, dough, bread, vinegar, like Lactobacillus acidophilus, L. lactis etc., Bacteria and Streptococcus lactis commonly called lactic acid bacteria (LAB) are probiotics which check the growth of pathogenic microbes in the stomach and other parts of the digestive tract. The LAB bacteria grows in milk and convert it into curd, thereby digesting the milk protein casein. A small amount of curd added to fresh milk as a starter or inoculum contains millions of Lactobacilli, which under suitable temperature (≤40oC) multiply and convert milk into curd. Curd is more nutritious than milk as it contains a number of organic acids and vitamins.
- Prebiotics are compounds in food (fibers) that induce the growth or activity of beneficial microorganisms.
- Probiotics are live microorganisms intended to provide health benefits when consumed, generally by improving or restoring the gut flora.
- Yogurt is produced by bacterial fermentation of milk, and lactic acid is produced as a by-product. Microorganisms such as *Streptococcus thermophilus* and Lactobacillus bulgaricus coagulate the milk protein and convert the lactose in the milk to lactic acid. The flavour in yogurt is due to acetaldehyde. Cheese is a dairy product produced in

95 | P a g e APPOLO STUDY CENTRE PH: 044-24339436, 42867555, 9840226187



a wide range of flavours, textures and is formed by coagulation of the milk protein, casein. During cheese production, milk is usually acidified and the enzyme rennet is added to cause coagulation. The solids are separated and pressed to form cheese. Most cheese are made with a starter bacteria, Lactococcus, Lactobacillus or Streptococcus.

- Paneer (cottage cheese) is fresh cheese common in South Asia, especially in India. It is made by curdling milk with lemon juice, vinegar and other edible acids. Large holes in Swiss cheese is due to the production of large amount of carbon-di-oxide by the bacterium Propionibacterium shermanii.
- The dough used in the preparation of idlis and dosas are fermented by the bacteria Leuconostoc mesenteroides whereas the dough used in bread making is fermented by Saccharomyces cerevisiae (Baker's Yeast). Fermentation of glucose mainly forms ethyl alcohol and carbon-di-oxide, which is responsible for leavening of dough. When leavened dough is baked, both carbon-di-oxide and ethyl alcohol evaporate making the bread porous and soft.

Single cell protein (SCP)

• Single cell protein refers to edible unicellular microorganisms like Spirulina. Protein extracts from pure or mixed cultures of algae, yeasts, fungi or bacteria may be used as ingredient or as a substitute for protein rich foods and is suitable for human consumption or as animal feed.

Microbes in industrial products

• Microbes are used to synthesize a number of products valuable to human beings. Products like beverages, antibiotics, organic acids, amino acids, vitamins, biofuels, single cell protein, enzymes, steroids, vaccines, pharmaceutical drugs, etc., are produced in industries. Production on a large scale requires growing microbes in very large vessels called fermentors. A fermentor (bioreactor) is a closed vessel with adequate arrangement for aeration, agitation, temperature, pH



control and drain or overflow vent to remove the waste biomass of cultured microorganisms along-with their products.

Antibiotic production

- Antibiotics are chemical substances produced by microorganisms which can kill or retard the growth of other disease causing microbes even in low concentration. Antibiotic means "against life". Antibiotics are used to treat diseases such as plague, meningitis, diphtheria, syphilis, leprosy, tuberculosis etc., Selman Waksman discovered Streptomycin and was the first to use the term "antibiotic" in 1943.
- While working on Staphylococci bacteria, Alexander Fleming observed a green mould growing in one of his unwashed culture plates around which Staphylococci could not grow. He found that it was due to a chemical produced by the mould and he named it as penicillin, which was the first antibiotic discovered by Alexander Fleming in 1926 (Fig. 9.1). Penicillin is produced by the fungi Penicillium notatum and Penicillium chrysogenum. It is bactericidal (antibiotics that kill bacteria) in action and inhibits the synthesis of the bacterial cell wall. Penicillin is also referred as the "queen of drugs" and its full potential as an effective antibiotic was established much later by Earnest Chain and Howard Florey when they treated the wounded soldiers in World War II with penicillin. Fleming, Chain and Florey were awarded the Nobel prize in 1945 for the discovery of penicillin.
- Antibiosis is the property of antibiotics to kill microorganisms.
- Broad-spectrum antibiotics act against a wide range of diseasecausing bacteria.
- Narrow-spectrum antibiotics are active against a selected group of bacterial types.

Hypersensitivity reaction is a major problem with the use of penicillin, resulting in nausea, vomiting, wheezing and ultimately cardiovascular collapse. To check the sensitivity reaction, doctors use a needle to prick the forearm of the patients to give a weak dose of penicillin. An itchy red region in the forearm is an indication that the patient is allergic to penicillin. This test is important before administration of penicillin to a

97 | P a g e APPOLO STUDY CENTRE PH: 044-24339436, 42867555, 9840226187

patient.



Tetracycline is a broad spectrum bacteriostatic antibiotic (antibiotics that limit the growth of bacteria) that inhibits microbial protein synthesis. Chlortetracycline is the first antibiotic of this group, of isolated from the cultures Streptomyces aureofaciens. Streptomycin is a broad spectrum antibiotic isolated from the actinomycetes, Streptomyces griseus. It is bactericidal against both gram positive and gram negative bacteria, especially against Mycobacterium tuberculosis. Antibiotics, such as erythromycin, chloromycetin, griseofulvin, neomycin, kenamycin, bacitracin, etc., are also isolated as microbial products.

Antibiotic resistance

• Antibiotic resistance occurs when bacteria develop the ability to defeat the drug designed to kill or inhibit their growth. It is one of the most acute threat to public health. Antibiotic resistance is accelerated by the misuse and over use of antibiotics, as well as poor infection prevention control. Antibiotics should be used only when prescribed by a certified health professional. When the bacteria become resistant, antibiotics cannot fight against them and the bacteria multiply. Narrow spectrum antibiotics are preferred over broad spectrum antibiotics. They effectively and accurately target specific pathogenic organisms and are less likely to cause resistance. "Superbug" is a term used to describe strains of bacteria that are resistant to the majority of antibiotics commonly used today.

Fermented beverages

 Microbes especially yeast is being used from time immemorial for the production of beverages like wine, beer, whisky, brandy and rum. Wine is among the oldest alcoholic beverages known and is produced by fermentation of fruit juice by yeast. Zymology is an applied science which deals with the biochemical process of fermentation and its practical uses. Saccharomyces cerevisiae commonly called brewer's yeast is used for fermenting malted cereals and fruit juices to produce various alcoholic beverages. Wine and beer are produced without distillation, whereas whisky, brandy and rum are obtained by fermentation and distillation.



• Oenology is the science and study of wine and wine making. Wine is made from the fermentation of grape juice. Grape juice is fermented by various strains of Saccharomyces cerevisiae into alcohol. Grape wine is of two types, red wine and white wine. For red wine, black grapes are used including skins and sometimes the stems also are used. In contrast white wine is produced only from the juice of either white or red grapes without their skin and stems. Beer is produced from germinated barley malt grain by Saccharomyces carlsbergensis or Saccharomyces cerevisiae. Rum is made from fermented sugarcane or molasses or directly from sugarcane juice by Saccharomyces cerevisiae. Whisky is a type of distilled alcoholic beverage made from fermented grain mash by Saccharomyces cerevisiae.

Alcohol content in various beverages

- Beer contains 3 to 5 percent of alcohol.
- Wine contains 9 to 14 percent alcohol. Wine coolers are made of wine mixed with carbonated water and flavourings. Wine coolers have about 4 to 6 percent alcohol.
- Distilled spirits such as whiskey, gin, scotch and vodka usually contain 35 to 50 percent alcohol.
- In some parts of South India, a traditional drink called pathaneer is obtained from fermenting sap of palms and coconut trees. A common source is tapping of unopened spadices of coconut. It is a refreshing drink, which on boiling produces jaggery or palm sugar. When pathaneer is left undisturbed for few hours it gets fermented to form toddy with the help of naturally occurring yeast, to form a beverage that contains 4 percent alcohol. After 24 hours toddy becomes unpalatable and is used for the production of vinegar.
- Saccharomyces cerevisiae is the major producer of ethanol (C2H5OH). It is used for industrial, laboratory and fuel purposes. So ethanol is referred to as industrial alcohol. Bacteria such as Zymomonas mobilis and Sarcina ventriculi are also involved in ethanol production. The principal substrates for the commercial production of industrial alcohol include molasses or corn, potatoes and wood wastes. The process of ethanol production starts by milling a feed stock followed by the addition of dilute or fungal amylase



(enzyme) from Aspergillus to break down the starch into fermentable sugars.

• Yeast is then added to convert the sugars to ethanol which is then distilled off to obtain ethanol which is upto 96 percent in concentration. The two most common type of biofuels in use today are ethanol and biodiesel, both of them represent the first generation of biofuel technology. Ethanol is often used as a fuel, mainly as a biofuel additive for gasoline.

Biodiesel is a fuel made from vegetable oils, fats or greases. Biodiesel fuel can be used in diesel engines without altering the engine. Pure biodiesel is non-toxic, biodegradable and produces lower level of air pollutants than petroleum-based diesel fuel. The Government of India approved the National Policy on Biofuels in December 2009 and identified Jatropha curcas as the most suitable oilseed for biodiesel production. Pongamia species is also a suitable choice for production of biodiesel.

Chemicals, enzymes and other bioactive molecules

- Microbes are not only used for commercial and industrial production of alcohol, but also used for production of chemicals like organic acids and enzymes. Examples of organic acid producers are Aspergillus niger for citric acid, Acetobacter aceti for acetic acid, Rhizopus oryzae for fumaric acid, Clostridium butyricum for butyric acid and Lactobacillus for lactic acid.
- Yeast (Saccharomyces cerevisiae) and bacteria are used for commercial production of enzymes. Lipases are used in detergent formulations and are used for removing oily stains from the laundry. Bottled juices are clarified by the use of pectinase, protease and cellulase. Rennet can also be used to separate milk into solid curds for cheese making. Streptokinase produced by the bacterium Streptococcus and genetically engineered Streptococci are used as "clot buster" for removing clots from the blood vessels of patients who have undergone myocardial infarction.
- Cyclosporin A, an immunosuppressant used in organ transplantation is produced from the fungus Trichoderma polysporum. It is also used

100 | P a g e APPOLO STUDY CENTRE PH: 044-24339436, 42867555, 9840226187



for its anti-inflammatory, anti-fungal and anti-parasitic properties. Statins produced by the yeast Monascus purpureus have been used to lower blood cholesterol levels. It acts by competitively inhibiting the enzyme responsible for the synthesis of cholesterol. Recombinant human insulin has been produced predominantly using E. coli and Saccharomyces cerevisiae for therapeutic use in human.

Microbes in sewage treatment and energy generation

Sewage is the waste generated every day in cities and towns containing human excreta. It contains large amounts of organic matter and microbes, which are pathogenic to humans and are biodegradable pollutants. Domestic waste consists of approximately 99 percent water, suspended solids and other soluble organic and inorganic substances. Sewage should not be discharged directly into natural water bodies like rivers and streams. Before disposal, sewage should be treated in sewage treatment plants to make it less polluting (Fig. 9.2). NTRE

Wastewater treatment

• The main objective of a wastewater treatment process is to reduce organic and inorganic components in wastewater to a level that it no longer supports microbial growth and to eliminate other potentially toxic materials. Microorganisms mainly bacteria and some protozoa play an essential part in the treatment of sewage to make it harmless. Sewage contains pathogenic bacteria. These bacteria must be destroyed in order to prevent the spread of diseases. Sewage treatment is usually performed in the following three stages.

Primary treatment

Primary treatment involves the physical removal of solid and particulate organic and inorganic materials from the sewage through filtration and sedimentation. Floating debris is removed by sequential filtration. Then the grit (soil and small pebbles) are removed by sedimentation. All solids that settle form the primary sludge and the supernatant forms the effluent. The effluent from the primary settling tank is taken for secondary treatment.



Secondary treatment or biological treatment

- The primary effluent is passed into large aeration tanks where it is constantly agitated mechanically and air is pumped into it. This allows vigorous growth of useful aerobic microbes into floc (masses of bacteria associated with fungal filaments to form mesh like structures). While growing, these microbes consume the major part of the organic matter in the effluent. This significantly reduces the BOD (Biochemical oxygen demand or Biological oxygen demand). BOD refers to the amount of the oxygen that would be consumed, if all the organic matter in one litre of water were oxidized by bacteria. The sewage water is treated till the BOD is reduced. The greater the BOD of the waste water more is its polluting potential.
- Once the BOD of sewage water is reduced significantly, the effluent is then passed into a settling tank where the bacterial "flocs" are allowed to sediment. This sediment is called activated sludge. A small part of activated sludge is pumped back into the aeration tank to serve as the inoculum. The remaining major part of the sludge is pumped into large tanks called anaerobic sludge digesters. Here, the bacteria which grow anaerobically, digest the bacteria and the fungi in the sludge. During this digestion, bacteria produce a mixture of gases such as methane, hydrogen sulphide and CO₂. These gases form biogas and can be used as a source of energy.

Tertiary treatment

• Tertiary treatment is the final process that improves the quality of the waste water before it is reused, recycled or released into natural water bodies. This treatment removes the remaining inorganic compounds and substances, such as nitrogen and phosphorus. UV is an ideal disinfectant for wastewater since it does not alter the water quality – except for inactivating microorganisms. UV is a chemical-free process that can completely replace the existing chlorination system and also inactivates chlorine-resistant microorganisms like Cryptosporidium and Giardia.



Act enforced by Government to conserve water bodies

- National river conservation plan (NRCP) was enacted in 1995 to improve the water quality of the rivers, which are the major fresh water resources in our country. This important assignment taken up under the NRCP includes,
 - ✓ To capture the raw sewage flowing into the river through open drains and divert them for treatment.
 - ✓ Setting up sewage treatment plants for treating the diverted sewage.
 - ✓ Construction of low cost sanitation toilets to prevent open defecation on river banks.
- The ministry for environment, forest and climate change has initiated the Ganga action plan and the Yamuna action plan to save the major rivers of the country. The Ganga action plan was launched on 14th January 1986. The main objective of the programme is to improve the water quality of River Ganges by interception, diversion and treatment of domestic sewage and to identify grossly polluting units to prevent pollution. The Yamuna Action Plan is a bilateral project between the Government of India and Japan. It was formally launched in April 1993. It was proposed to build large number of sewage treatment plants to discharge treated wastewater into the rivers.

Microbial fuel cell (MFC)

• A microbial fuel cell is a bio-electrochemical system that drives an electric current by using bacteria and mimicking bacterial interaction found in nature (Fig. 9.3). Microbial fuel cells work by allowing bacteria to oxidize and reduce organic molecules. Bacterial respiration is basically one big redox reaction in which electrons are being moved around. A MFC consists of an anode and a cathode separated by a proton exchange membrane. Microbes at the anode oxidize the organic fuel generating protons which pass through the membrane to the cathode and the electrons pass through the anode to the external circuit to generate current.



Microbes in the production of biogas (Gobar gas)

- Biogas is a mixture of different gases produced by the breakdown of organic matter in the absence of oxygen. Biogas can be produced from raw materials such as agricultural wastes, manure, municipal wastes, plant material, sewage, food waste, etc., Biogas is produced under anaerobic condition, when the organic materials are converted through microbiological reactions into gas and organic fertilizer. Biogas primarily consists of methane (63 percent), along with CO2 and hydrogen. Methane producing bacteria are called methanogens and one such common bacterium is Methano bacterium. Biogas is devoid of smell and burns with a blue flame without smoke. The Methanogens are also present in anaerobic sludge and rumen of cattle.
- In rumen, these bacteria help in the breakdown of cellulose. The excreta of cattle called dung is commonly called "Gobar". Gobar gas is generated by the anaerobic decomposition of cattle dung. It consists of methane, CO2 with some hydrogen, nitrogen and other gases in trace amounts.
- In a biogas plant, anaerobic digestion is carried out in an air tight cylindrical tank known as digester (Fig. 9.4). It is made up of concrete bricks and cement or steel. Bio-wastes are collected and slurry of dung is fed into this digester. It has a side opening into which organic materials for digestion are incorporated for microbial activity. Anaerobic digestion is accomplished in three stages: solubilisation, acidogenesis and methanogenisis. The outlet is connected to a pipe to supply biogas. The slurry is drained through another outlet and is used as fertilizer. Biogas is used for cooking and lighting. The technology of biogas production was developed in India mainly due to the efforts of Indian Agricultural Research Institute (IARI) and Khadi and Village Industries Commission (KVIC).

Microbes as bio control agents and bio fertilisers

• Large scale application of chemical insecticides and pesticides have a deleterious effect on the health of human beings and pollute our environment. Bio control is a method of controlling pest by use of microbes such as fungi, bacteria, viruses or by naturally occurring



substances derived from plants and animals. The use of a microbes or other biological agents to control a specific pest is called a biopesticide. Biopesticides are used to control insect pests. The lady bird beetle and dragonflies are useful to control aphids and mosquito larvae respectively. Bacillus thuringiensis is a soil dwelling bacterium which is commonly used as a biopesticide and contains a toxin called cry toxin (Fig. 9.5). Scientists have introduced this toxin producing genes into plants and have raised genetically engineered insect resistant plants. E.g. Bt-cotton.

- During sporulation Bacillus thuringiensis produces crystal proteins called Delta-endotoxin which is encoded by cry genes. Delta-endotoxins have specific activities against the insects of the orders Lepidoptera, Diptera, Coleoptera and Hymenoptera. When the insects ingest the toxin crystals their alkaline digestive tract denatures the insoluble crystals making them soluble. The cry toxin then gets inserted into the gut cell membrance and paralyzes the digestive tract. The insect then stops eating and starves to death.
- Weedicides are substances, which destroy weeds without harming the useful plants. Bio weedicides are compounds and secondary metabolites derived from microbes such as fungi, bacteria or protozoa. The first bio herbicide developed in 1981 was a Mycoheribicide derived from the fungus Phytophthora palmivora. It controls the growth of strangler vine in citrus crops. Trichoderma species are free living fungi that are very common in the root ecosystem. They are effective bio control agents for several plant pathogens. Buculoviruses are pathogens that attack insects and other arthropods. The genus Nucleopolyhedrovirus is used as a biocontrol agent. These viruses are species specific and have narrow spectrum insecticidal applications.

Biofertilisers

• Biofertilisers are formulation of living microorganisms that enrich the nutrient quality of the soil. They increase physico – chemical properties of soils such as soil structure, texture, water holding capacity, cation exchange capacity and pH by providing several nutrients and sufficient organic matter. The main sources of



biofertilisers are bacteria, fungi and cyanobacteria. Rhizobium is a classical example for symbiotic nitrogen fixing bacteria. This bacterium infects the root nodules of leguminous plants and fixes atmospheric nitrogen into organic forms. Azospirillum and Azotobacter are free living bacteria that fix atmospheric nitrogen and enrich the nitrogen content of soil.

- A symbiotic association between a fungus and the roots of the plants is called mycorrhiza. The fungal symbiont in these associations absorbs the phosphorus from soil and transfers to the plant. Plants having such association show other benefits such as resistance to root-borne pathogens, tolerance to salinity, drought, enhances plant growth and developments. For example, many members of the genus Glomus form mycorrhiza. Cyanobacteria (or) blue green algae (BGA) are prokaryotic free-living organisms which can fix nitrogen. Oscillatoria, Nostoc, Anabaena, Tolypothrix are well known nitrogen fixing cyanobacteria. Their importance is realized in the water logged paddy fields where Cyanobacteria multiply and fix molecular nitrogen. Cyanobacteria secrete growth promoting substances like indole-3-acetic acid, indole-3- butyric acid, naphthalene acetic acid, amino acids, proteins, vitamins which promotes plant growth and production.
- Bio fertilisers are commonly used in organic farming methods. Organic farming is a technique, which involves cultivation of plants and rearing of animals in natural ways. This process involves the use of biological materials, avoiding synthetic substances to maintain soil fertility and ecological balance thereby minimizing pollution and wastage.

Key features of organic farming

- Protecting soil quality using organic materials and encouraging biological activity.
- Indirect provision of crop nutrients using soil microorganisms.
- Nitrogen fixation in soils using legumes.



Weed and pest control based on methods like crop rotation, biological diversity, natural predators, organic manures and suitable chemical, thermal and biological interventions.

Bioremediation

or genetically The use naturally occurring engineered of degrade pollutants microorganisms to reduce or is called bioremediation. Bioremediation is less expensive and more sustainable than other remediation's available. It is grouped into in situ bioremediation (treatment of contaminated soil or water in the site) and ex situ bioremediation (treatment of contaminated soil or water that is removed from the site and treated).

Microorganisms involved in bioremediation

- Aerobic microbes degrade the pollutants in the presence of oxygen. They mainly degrade pesticides and hydrocarbons. Pseudomonas putida is a genetically engineered microorganism (GEM). Ananda Mohan Chakrabarty obtained patent for this recombinant bacterial strain. It is multi-plasmid hydrocarbon-degrading bacterium which can digest the hydrocarbons in the oil spills (Fig. 9.6).
- Nitrosomonas europaea is also capable of degrading benzene and a variety of halogenated organic compounds including trichloroethylene and vinyl chloride. Ideonella sakaiensis is currently tried for recycling of PET plastics (Fig. 9.7). These bacteria use PETase and MHETase enzymes to breakdown PET plastic into terephthalic acid and ethylene glycol.
- Anaerobic microbes degrade the pollutants in the absence of oxygen. Dechloromonas aromatica has the ability to degrade benzene anaerobically and to oxidize toluene and xylene. Phanerochaete chrysosporium an anaerobic fungus exhibits strong potential for bioremediation of pesticides, polyaromatic hydrocarbons, dyes, trinitrotoluene, cyanides, carbon tetrachloride, etc., Dehalococcoides species are responsible for anaerobic bioremediation of toxic trichloroethene to non-toxic ethane. Pestalotiopsis microspora is a species of endophytic fungus capable of breaking down and digesting



polyurethane. This makes the fungus a potential candidate for bioremediation projects involving large quantities of plastics.

• Ideonella sakaiensis adhere to PET film PET Bottle Eats the terephthalic acid & ethylene glycol Breaks down PET into terephthalic acid & ethylene glycol Fig. 9.7 Actions of Ideonella sakaiensis

Summary

 All microbes are not pathogenic, many of them are beneficial to human beings. We use microbes and their derived products almost every day. Lactic acid bacteria convert milk into curd. Saccharomyces cerevisiae (yeast) is used in bread making. Idly and dosa are made from dough fermented by microbes. Bacteria and fungi are used in cheese making. Industrial products like lactic acid, acetic acid and alcohol are produced by microbes. Antibiotics are produced from useful microbes to kill the disease causing harmful microbes. For more than a hundred years, microbes are being used to treat sewage by the process of activated sludge formation. Bio-gas produced by microbes is used as a source of energy in rural areas. Microbes are also used as bio-control agents to avoid the use of toxic pesticides. Now a days chemical fertilisers are gradually replaced by biofertilisers. In bio-remediation naturally occurring or genetically engineered microorganisms are used to reduce or degrade pollutants.